CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 101, 2008

A

Abbott, V. A., 784
Abel, C. A., 168, 1406
Abellán, J., 499, 1685
Abney, M. R., 23, 1238
Adachi-Hagimori, T., 1510
Adalla, C. B., 575
Adamczyk, J. J., Jr, 1594
Adaniya, K., 1568
Ahmad, Munir, 472, 1667, 1927
Ahmad, Mushtaq, 1667
Ahn, Y.-J., 139, 1960
Akbar, W., 1903
Akhtar, Z.-R., 1415
Alavi, S., 989
Albright, R. L., 1095
Alfaro, F., 126
All, J. N., 1471, 1950
Allen, C. T., 1729
Allen, M. L., 265
Amarasekare, K. G., 1798
Andow, D. A., 1406
Anfora, G., 444
Angeli, G., 444
Anhold, J. A., 1253
Anilkumar, K. J., 1421
Appel, A. G., 151
Arakaki, N., 1568
Aranda-Herrera, E., 769
Armstrong, J. S., 399, 742
Armstrong, K. F., 523
Arthurs, S. P., 288, 1540
Asahara, M., 1704
Ascunce, M. S., 1443
Aslam, M., 1667, 1676
Assefa, Y., 686
Athanassiou, C. G., 976
Atkinson, E. B., 838
Atlihan, R., 325
Aupinel, P., 1737

B

Bacheler, J., 1950
Backus, E. A., 1183
Bai, L.-Y., 1888
Balasubramani, V., 1911
Baldessari, M., 444
Baldwin, J., 492
Baldwin, R. W., 1384
Ball, S. L., 523
Bartlett, J. S., 61
Battany, M. C., 699
Baxendale, F. P., 533
Behle, R. W., 797, 1856
Bell, C. E., 1624
Benjamin, T., 1779
Bennett, G. W., 894
Bensen, T. A., 1526
Bernklau, E. J., 341
Berthomieu, A., 484
Berticat, C., 484
Bertolaccini, I., 1792
Bettiga, L. J., 1515
Beuzelin, J. M., 1903
Bextine, B., 1122
Bible, J. B., 1584
Bielza, P., 499, 1685
Bishop, B. L., 1158
Bjostad, L. B., 341
Blanco, C. A., 168, 1406
Blodgett, S., 309
Bloem, K. A., 314
Bloem, S., 314
Blua, M. J., 1122
Boerma, H. R., 1471
Boetel, M. A., 1760
Boina, D. R., 989
Boiteau, G., 56, 1575
Bommireddy, P. L., 1244
Bonte, M., 1127
Borden, J. H., 1266
Borgemeister, C., 90
Bosch, J., 647, 660, 1088
Bosque-Pérez, N. A., 1606
Bostanian, N. J., 145
Bosu, P. P., 944
Boukouvala, M. C., 976
Boulton, A., 1331
Bourham, M. A., 302
Bradley, J. R., 1950
Bradshaw, J. D., 801, 1211
Braxton, L. B., 1950
Brewer, G. J., 969
Brewer, M. J., 278
Brown, G. R., 546
Brown, J. J., 199
Brown, S. L., 631
Bruck, D. J., 251
Brunner, J. F., 1651
Buckelew, L., 174
Buczkowski, G., 894
Buntin, G. D., 631, 742, 1455
Bures, B. L., 302
Burkness, E. C., 461
Burks, C. S., 1633
Burks, J. ", 555
Burley, L. M., 1081
Burrack, H. J., 750
Buss, E. A., 1847
Büyükgüzel, K., 777
Byers, J. A., 720

C

Caballero, P., 42
Caboni, P., 1075
Cabras, P., 1075
Cabrera-La Rosa, J. C., 1105
Calderone, N. W., 668
Callcott, A.-M. A., 1158
Calvin, D., 1614
Campan, E., 584
Campbell, J. F., 631, 1719
Cantrell, C. L., 265
Cardona, C., 564
Carpenter, J. E., 314
Carraher, C., 1944
Carrière, Y., 504, 927
Carroll, M. W., 1614
Casanoves, F., 1779
Castle, S. J., 174, 226
Catchot, A., 1594
Cavichioli, R. R., 1114
Cermak, S. C., 797
Chabi-Olaye, A., 90
Chaney, W. E., 1526
Chang, G. C., 74
Chang, J., 933
Chang, L. H., 1743
Chapman, N. C., 272
Cheikh, H. B., 484
Cheikh, R. B., 484
Chen, B. L., 637
Chen, H., 182
Chen, J., 265
Chen, Jianchi, 1105
Chen, Jin, 515, 933
Chen, M., 438
Chen, W.-L., 1760
Chen, Y. H., 575
Chen, Y. L., 1743
Cheng, J.-A., 1415
Chi, H., 325
Chi, Z.-J., 360
Chintzoglou, G. J., 976
Cho, H. S., 1134
Choe, D.-H., 1397
Choi, D.-R., 1960
Chong, J.-H., 1798
Chouinard, G., 1838
Chouvenc, T., 885
Chuang, Y. C., 1743
Chuang, Y.-Y., 352
Cicciarelli, R. P., 668
Civerolo, E. L., 1105
Clancy, K. M., 1253
Clark, A., 1443
Clarke, S. R., 1821

CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 101, 2008

A

Abbott, V. A., 784
Abel, C. A., 168, 1406
Abellán, J., 499, 1685
Abney, M. R., 23, 1238
Adachi-Hagimori, T., 1510
Adalla, C. B., 575
Adamczyk, J. J., Jr, 1594
Adaniya, K., 1568
Ahmad, Munir, 472, 1667, 1927
Ahmad, Mushtaq, 1667
Ahn, Y.-J., 139, 1960
Akbar, W., 1903
Akhtar, Z.-R., 1415
Alavi, S., 989
Albright, R. L., 1095
Alfaro, F., 126
All, J. N., 1471, 1950
Allen, C. T., 1729
Allen, M. L., 265
Amarasekare, K. G., 1798
Andow, D. A., 1406
Anfora, G., 444
Angeli, G., 444
Anhold, J. A., 1253
Anilkumar, K. J., 1421
Appel, A. G., 151
Arakaki, N., 1568
Aranda-Herrera, E., 769
Armstrong, J. S., 399, 742
Armstrong, K. F., 523
Arthurs, S. P., 288, 1540
Asahara, M., 1704
Ascunce, M. S., 1443
Aslam, M., 1667, 1676
Assefa, Y., 686
Athanassiou, C. G., 976
Atkinson, E. B., 838
Atlihan, R., 325
Aupinel, P., 1737

B

Bacheler, J., 1950
Backus, E. A., 1183
Bai, L.-Y., 1888
Balasubramani, V., 1911
Baldessari, M., 444
Baldwin, J., 492
Baldwin, R. W., 1384
Ball, S. L., 523
Bartlett, J. S., 61
Battany, M. C., 699
Baxendale, F. P., 533
Behle, R. W., 797, 1856
Bell, C. E., 1624
Benjamin, T., 1779
Bennett, G. W., 894
Bensen, T. A., 1526
Bernklau, E. J., 341
Berthomieu, A., 484
Berticat, C., 484
Bertolaccini, I., 1792
Bettiga, L. J., 1515
Beuzelin, J. M., 1903
Bextine, B., 1122
Bible, J. B., 1584
Bielza, P., 499, 1685
Bishop, B. L., 1158
Bjostad, L. B., 341
Blanco, C. A., 168, 1406
Blodgett, S., 309
Bloem, K. A., 314
Bloem, S., 314
Blua, M. J., 1122
Boerma, H. R., 1471
Boetel, M. A., 1760
Boina, D. R., 989
Boiteau, G., 56, 1575
Bommireddy, P. L., 1244
Bonte, M., 1127
Borden, J. H., 1266
Borgemeister, C., 90
Bosch, J., 647, 660, 1088
Bosque-Pérez, N. A., 1606
Bostanian, N. J., 145
Bosu, P. P., 944
Boukouvala, M. C., 976
Boulton, A., 1331
Bourham, M. A., 302
Bradley, J. R., 1950
Bradshaw, J. D., 801, 1211
Braxton, L. B., 1950
Brewer, G. J., 969
Brewer, M. J., 278
Brown, G. R., 546
Brown, J. J., 199
Brown, S. L., 631
Bruck, D. J., 251
Brunner, J. F., 1651
Buckelew, L., 174
Buczkowski, G., 894
Buntin, G. D., 631, 742, 1455
Bures, B. L., 302
Burkness, E. C., 461
Burks, C. S., 1633
Burks, J. ", 555
Burley, L. M., 1081
Burrack, H. J., 750
Buss, E. A., 1847
Büyükgüzel, K., 777
Byers, J. A., 720

C

Caballero, P., 42
Caboni, P., 1075
Cabras, P., 1075
Cabrera-La Rosa, J. C., 1105
Calderone, N. W., 668
Callcott, A.-M. A., 1158
Calvin, D., 1614
Campan, E., 584
Campbell, J. F., 631, 1719
Cantrell, C. L., 265
Cardona, C., 564
Carpenter, J. E., 314
Carraher, C., 1944
Carrière, Y., 504, 927
Carroll, M. W., 1614
Casanoves, F., 1779
Castle, S. J., 174, 226
Catchot, A., 1594
Cavichioli, R. R., 1114
Cermak, S. C., 797
Chabi-Olaye, A., 90
Chaney, W. E., 1526
Chang, G. C., 74
Chang, J., 933
Chang, L. H., 1743
Chapman, N. C., 272
Cheikh, H. B., 484
Cheikh, R. B., 484
Chen, B. L., 637
Chen, H., 182
Chen, J., 265
Chen, Jianchi, 1105
Chen, Jin, 515, 933
Chen, M., 438
Chen, W.-L., 1760
Chen, Y. H., 575
Chen, Y. L., 1743
Cheng, J.-A., 1415
Chi, H., 325
Chi, Z.-J., 360
Chintzoglou, G. J., 976
Cho, H. S., 1134
Choe, D.-H., 1397
Choi, D.-R., 1960
Chong, J.-H., 1798
Chouinard, G., 1838
Chouvenc, T., 885
Chuang, Y. C., 1743
Chuang, Y.-Y., 352
Cicciarelli, R. P., 668
Civerolo, E. L., 1105
Clancy, K. M., 1253
Clark, A., 1443
Clarke, S. R., 1821

Clodius, M., 365
Colby, D., 1062
Coler, R., 759
Conlong, D. E., 686
Cook, D. R., 492
Cooper, M. L., 699
Copenhaver, K., 1614
Cornelius, M. L., 1367
Cossé, A. A., 1856
Costamagna, A. C., 1488
Costello, M. J., 1287
Costello, S. L., 409
Cox-Foster, D. L., 256
Crickmore, N., 1658, 1911, 1927
Croft, B. J., 17
Crowder, D. W., 927
Cui, F., 1897
Cullen, E. M., 391

D

Daane, K. M., 699, 1515
Dai, D. J., 515
Damos, P. T., 1557
Danka, R., 654
Davis, J. A., 9
Davis, R. W., 902
Davis, T., 1643
De Clercq, P., 1127
De Cristofaro, A., 444
Degain, B. A., 927
Degomez, T. E., 1253
de la Rosa, F., 1540
Demchak, K., 850
Dennehy, T. J., 504, 927
Dennis, S. O., 1158
Denno, R. F., 959
Dhileepan, K., 17
Dickens, J. C., 859
Dively, G. P., 959
Dixon, P. L., 822
Dobson, S., 1944
Doerr, M., 1651
Domínguez, J., 126
Domínguez-Ruiz, J., 1325
Donahue, K. M., 251
Dong, P., 1711
Donohue, K. V., 302
Dorhout, D. L., 404
Dorn, S., 81, 314, 1341
Dosdall, L. M., 159
Doskocil, J. P., 1624
Dou, W., 1711
Downes, S., 909
Drummond, F. A., 334
Dubois, F., 584
Duerr, D. A., 107
Dugravot, S., 584
Duke, S. O., 265
Dunlap, C., 1856
Durmuş, Y., 777

E

Eaton, J. A., 199
Edwards, D. L., 251
Edwards, R., 1944
Eguaras, M., 1075
Eichenseer, H., 555
Eickhoff, T. E., 533
Eigenbrode, S. D., 1606
Eliopoulos, P. A., 982
Ellers-Kirk, C., 504, 927
Elliott, M. L., 885
El-Sayed, A. M., 720
Epsky, N. D., 1798
Ernst, J. A., 1443
Espino, L., 216, 1233
Espinoza, J. L., 190

F

Fandohan, P., 1967
Fang, Q., 1415
Fell, R. D., 1081
Fernández, E., 499, 1685
Finley, J., 1057
Fisher, J., 1856
Flanders, K., 742
Flint, M. L., 829
Flores, D., 1478
Floris, I., 1075
Foil, L. D., 1034, 1062
Follett, P. A., 710, 1068
Forsline, P. L., 206
Foster, J. E., 533
Fraedrich, S. W., 1276
Francese, J. A., 1831
Fraser, I., 1831
French, J. V., 1478
Fridgen, J., 1614
Fry, H.R.C., 822
Fu, A.A.A., 769
Fuentes-Contreras, E., 190
Fuester, R. W., 1140

G

Gage, B., 365
Gao, C. F., 515
Gassmann, A. J., 504
Gaul, S., 118
Gera, A., 1337
Germinara, G. S., 444
Getchell, A. I., 1022
Getu, E., 50
Ghadamyari, M., 1432
Giguère, M.-A., 728
Giordanengo, P., 584
Glaser, J. A., 1614
Gnonlonfin, G.J.B., 1967
Gorden, J. M., 451
Gore, J., 742, 1594
Gould, F., 1406
Grafton-Cardwell, E. E., 451, 1041
Grávalos, C., 499, 1685
Greenberg, S. M., 399
Greenplate, J. T., 546
Grenkow, L. F., 1811
Groves, R. L., 1041, 1105
Gu, H.-N., 384
Guedes, R.N.C., 1974
Gunawan, G., 631
Guo, B., 1455
Guy, S. O., 1606

H

Hackenberger, B. K., 1204
Haley, S. D., 569
Hall, D. G., 1478
Hallman, G. J., 716, 1051
Hamilton, G. C., 1439
Han, L. Z., 1691
Hanavan, R. P., 1606
Hannon, E. R., 199
Hanson, B. K., 810
Hanula, J. L., 1146, 1276
Haraguchi, D., 1198
Harpold, V. S., 927
Harris, M. K., 769, 1584
Harrison, H. F., Jr, 1871
Hartley, D., 1934
Hartman, G. L., 1228
Hayes, C. J., 1253
He, X. Z., 1771
Head, M., 623
Headrick, H. L., 288
Hell, K., 1967
Hellmich, R. L., 1614
Heng-Moss, T. M., 533
Henson, R. A., 810
Heraty, J. M., 1771
Herbert, K., 1934
Herdina,, 1934
Hernández, G., 1406
Herrman, T., 998
Hertlein, M., 759
Herzog, G. A., 23
Heuberger, S., 504
Hibbard, B. E., 797, 1547
Higa, M., 1568
Higbee, B. S., 1633
Hight, S. D., 314
Higo, H. A., 784
Hildago, E., 1779
Hill, G., 1944
Hill, J. H., 1211
Hill, M. P., 99
Hinomoto, N., 1167
Hitchner, E. M., 859
Hoddle, C. D., 692, 1310
Hoddle, M. S., 692, 1310
Hoffmann, A., 1934
Hogsette, J. A., 1034
Hokama, Y., 1568
Holbrook, C. C., 631
Hong, X.-Y., 360
Horsfield, A., 334
Hou, B., 736

Hou, M. L., 1691
Hou, R. F., 352
Huang, C.-H., 1415
Huang, F., 492, 1903
Huang, X.-Y., 1888
Hubert, J., 1028
Huckaba, R. M., 1950
Hunt, T. E., 1614
Huntzinger, C. I., 660, 1088
Husseneder, C., 1062
Hutchison, W. D., 461

I

Imamura, T., 1006
Imamura, Y., 880
Ioriatti, C., 444
Irigoyen, J., 132
Irvine, R. B., 1811
Isbell, T. A., 797

J

Jackman, J. A., 1584
Jackson, B. C., 1122
Jackson, D. M., 1871
Jackson, R., 168, 742, 1594
Jacobi, W. R., 409
Jahn, G. C., 575
James, R. R., 660, 674, 1088
James, S. S., 1158
Jamieson, L. E., 1295
Jarić-Perkušić, D., 1204
Jasrotia, P., 1238
Jayas, D. S., 1011
Jiang, C., 546
Jilani, G., 1676
Johnson, K. D., 801
Johnson, K. N., 17
Johnson, M. W., 1105
Jones, V. P., 1651
Jong, D. Y., 838
Jürgens, A., 720

K

Kamble, S. T., 902, 1373
Kang, J. S., 1134
Kaplan, I., 959
Katsumata, N., 880
Kavallieratos, N. G., 976
Kawamitsu, H., 1568
Keddie, B. A., 159
Kemp, W. P., 647, 660, 1088
Kennedy, G. G., 1238
Kim, J. I., 1134
Kim, J. N., 139
Kim, J.-R., 1960
Kim, K. S., 1729
Kim, Y., 36
Kim, Y. H., 1134
Kishita, M., 1568
Kittelson, N. T., 199
Klavons, J. A., 1697
Klein, M. G., 1158
Knodel, J. J., 810
Knutson, A. E., 1501
Kobayashi, A., 1568
Koehler, P. G., 1384, 1389, 1881
Kohama, T., 1198
Kono, Y., 1432
Kopittke, R. A., 61
Koppenhöfer, A. M., 1856
Kostromytska, O. S., 1847
Kotsubo, Y., 1167
Kraiss, H., 391
Krakowsky, M. D., 1455
Kring, T. J., 23
Kudlikova-Krizkova, I., 1028
Kuhar, T. P., 859
Kumano, N., 1198
Kundu, S. C., 1176
Kwon, B., 36
Kwon, M., 1134

L

Lacasa, A., 1864
Lacey, L. A., 288, 1540
Ladurner, E., 647
Lafontaine, J. P., 1266
Lambkin, T. A., 61
Lamothe, S., 1838
Lance, D. R., 1831
Larentzaki, E., 1317
Larson, Z., 998
Lasa, R., 42
Lavandero, B., 190
Lax, A. R., 1367
Lee, B., 623
Lee, D. R., 492
Lee, E.-J., 1960
Lee, H.-C., 1152
Lee, J. C., 829
Lee, J. E., 451
Lee, R. D., 631
Lee, S., 1134
Lee, S.-G., 139
Lee, S. H., 1134
Lee, S.-H., 592
Lee, Y.-S., 139
Legendre, B. L., 237
Lemoyne, P., 118
Leonard, B. R., 1244, 1950
Leonard, B. Rogers, 492, 742
Leonard, R., 168
Leopold, R. A., 1760
Le Roux, V., 584
Leskey, T. C., 1302
Leung, H., 575
Li, A. Y., 1697
Li, W. H., 515
Lim, J., 272
Lin, Y., 182
Liu, F., 933
Liu, J.-L., 384
Liu, P. L., 1691
Liu, X.-J., 360
Liu, Y., 1805
Liu, Y.-B., 1515, 1786
Loeb, G. M., 850
Lopes, J.R.S., 1114
Lopez, J., 742
Lopez, J. D., Jr, 168
Loughner, R. L., 850
Lu, Y., 1805

M

Ma, C. Y., 515
Macías, J., 1821
MacKenzie, K., 118
Mahendran, B., 1176
Maheswaran, M., 575
Mahon, D., 623
Mahon, R. J., 909
Maier, C. T., 430
Maini, S., 647
Mankin, R. W., 838
Manning, L. M., 720
Mannion, C. M., 1798
Marek, S. M., 1624
Marucci, R. C., 1114
Mastro, V. C., 1831
Mauchline, N., 1944
Mayfield, A. E., III, 1276
Mbungu, N. T., 56
McAnally, L. R., 1158
McClure, M., 728
McCornack, B. P., 1488
McElfresh, J. S., 769
McKay, A., 1934
McKenna, C., 1944
McMillin, J. D., 1253
Meagher, R. L., 742
Meikle, W. G., 1719
Melathopoulos, A. P., 1095
Meléndez-Ramirez, V., 132
Meng, F., 933
Merdić, E., 1204
Merrill, S. C., 541, 955
Metz, B. N., 1749
Michels, G. J., 1584
Midgley, J. M., 99
Miles, J. W., 564
Millar, J. G., 769, 1771
Miller, D. R., 107
Miller, H. R., 541
Miller, J. A., 1697
Miller, M. F., 564
Miller, R. S., 309
Miller, T. A., 1519
Mitchell, A., 686
Mitchell, V. J., 720
Miura, K., 1510
Miyanoshita, A., 1006
Mizuno, H., 1432
Mo, J., 1331
Moar, W. J., 546, 1421
Mody, K., 81, 1341
Mogi, N., 1568

Montez, G., 1041
Montoya, J., 564
Moore, S. H., 492
Moreira, J. A., 769
Moreno, B., 1821
Mottern, J. L., 1771
Moyseenko, J. J., 1158
Mpinga, M., 114
Muegge, M. A., 1501
Muirhead, K. A., 686
Mullens, B. A., 637
Mullins, J. W., 546
Mulungu, L. S., 114
Mundal, K. D., 969
Munro, S., 1331
Munzbergova, Z., 1028
Murata, M., 1006
Mwatawala, M. W., 114
Myers, C. T., 206
Myers, H. M., 480

N

Nadeau, J. L., 784
Nagayama, A., 1568
Nagoshi, R. N., 742
Nakama, S., 1568
Nalli, A., 1034
Nanos, G. D., 866
Nansen, C., 1719
Naranjo, S. E., 226, 1501
Nault, B. A., 1317
Navarro-Llopis, V., 126, 1325
Neerdaels, N. B., 1238
Negrón, J. F., 409
Nelson, E. H., 699
Neto, A. M., 759
Neumann, G., 1533
Neven, L. G., 295
Newcomb, R. D., 1944
Ni, X., 631, 1455
Nie, X., 728
Nielsen, A. L., 1439
Nigg, H. N., 1443
Nolting, S., 1950
Noma, T., 278
Noor-Ul-Ane, M., 1658
Nunez, H., 769
Núñez-Pérez, E., 1792

O

Oldroyd, B. P., 272
Oliver, J. B., 1158, 1831
Olmer, K. J., 1547
Olsen, K. M., 909
Olson, B. E., 309
Olson, D. L., 810
O'Neal, M. E., 801
O'Neill, K. M., 309
Ophel-Keller, K., 1934
Opit, G. P., 605, 616, 1974
Oppenhuizen, M. E., 546
Orr, D. B., 421
Osakabe, Mh., 1167, 1704
Osborn, J. A., 546
Osbrink, W.L.A., 1367
Ottea, J. A., 1903

P

Paine, T. D., 1771
Palchamy, K., 575
Palma, G., 132
Pankiw, T., 1749
Panoussakis, E. C., 976
Papachristos, D. P., 866
Papadopoulos, N. T., 866
Park, I.-K., 139
Parman, V., 1349
Parrott, W. A., 1471
Pasteur, N., 484
Pathan, A. K., 1676
Payne, G., 168
Peairs, F. B., 541, 569, 955
Pelletier, Y., 728
Peng, Y. F., 1691
Pereira, R. M., 1384, 1389, 1881
Perera, O. P., 1406
Pernal, S. F., 1095
Perring, T. M., 1519
Peteinatos, G. G., 976
Petrou, S. S., 976
Pfeiffer, D. G., 859
Pfiester, M., 1389
Philip, M. M., 421
Phillips, T. W., 1051, 1719
Picka, J. D., 1575
Piñero, J. C., 759, 1302
Pitts-Singer, T. L., 674
Pivonia, S., 1337
Plate, J., 438, 1317
Poulton, J., 1944
Powell, K., 1934
Prabhaker, N., 174
Prado, J., 1779
Price, D. K., 1068
Primo, J., 126, 1325
Prokopy, R. J., 1302
Pureswaran, D. S., 1266
Pusztai-Carey, M., 1421
Puterka, G. J., 569

Q

Qiao, C., 1897
Qiao, F., 915
Quezada-Euán, J.J.G., 132
Quinto, V., 499, 1685
Quiring, D. T., 822

R

Rabaglia, R. J., 1276
Racette, G., 145
Radcliffe, E. B., 9
Ragsdale, D. W., 1488
Ramankutty, P., 1295
Ramaseshiah, G., 1140
Ramirez, J. L., 1519
Ramírez, C. C., 190
Randolph, T. L., 541, 569, 955
Razaq, M., 1676
Reagan, T. E., 237, 1903
Reay-Jones, F.P.F., 237
Rebek, E., 1643
Rector, B. G., 1455
Reding, M. E., 1158
Reed, D. A., 1771
Reggiori, F., 444
Reid, R., 623
Reinert, J. A., 1624
Reissig, W. H., 206
Rejón, M., 132
Ren, Y., 623
Rice, M. E., 404, 801, 1211
Rice, S. J., 61
Richman, D., 480
Ridley, A. W., 17
Rinderer, T. E., 1756
Roberts, P. M., 23
Robillard, S. M., 451
Robinson, J.R.C., 1600
Roe, R. M., 302
Ross, D. W., 1826
Royer, T. A., 1624
Rozelle, S., 915
Ruberson, J. R., 23, 631
Rudolph, J. B., 541, 569
Ruschke, L. G., 546
Rust, M. K., 699, 873, 1360, 1397
Ryall, K. L., 822

S

Saacke, R. G., 1081
Saavedra, H. O., 365, 375
Sadof, C., 1779
Sadoyama, Y., 1568
Saeed, S., 1658
Sagili, R. R., 1749
Saha, M., 1176
Saleem, M. A., 472, 1667, 1676
Sallam, M.N.S., 334
Sammataro, D., 1057
Sánchez, J. A., 1864
Sanchis, J., 126, 1325
Sandanayaka, M., 1944
Sandanayaka, W.R.M., 1183
Sangha, J. S., 575
Sansone, C., 742
Sappington, T. W., 1614, 1729
Saran, R. K., 873, 1360, 1373
Sarfraz, R. M., 159
Sarvary, M. A., 314
Satta, A., 1075
Savopoulou-Soultani, M., 1557
Sawyer, A. J., 1831
Sayyed, A. H., 472, 1658, 1667, 1676, 1911, 1927
Scherer, C. W., 894
Schiffer, M., 1934

Schloemann, S., 850
Schotzko, D. J., 1606
Schulthess, F., 90
Schultz, P. B., 859
Schultz, R. J., 334
Sebaugh, J. L., 546
Sétamou, M., 1478
Seybold, S. J., 829
Seyoum, E., 50
Shappley, Z. W., 546
Shearer, P. W., 1439
Shelton, A. M., 438, 1317
Shen, J. L., 515, 933
Shields, E. J., 1533
Shin, S.-C., 139
Shinzato, T., 1568
Showler, A. T., 68, 1600
Siame, A. B., 1967
Siebert, M. W., 1950
Siegfried, B. D., 168
Sime, K. R., 699
Sisterson, M. S., 1, 1041
Sivasupramaniam, S., 546
Smith, H. A., 1526
Smith, J. A., 1881
Smith, L. M., II, 151
Smith, M. T., 838
Smitley, D., 1643
Snook, M. E., 1455
Snyder, W. E., 74
Sobotnik, J., 1028
Soroka, J. J., 1811
Sotelo, G., 564
Sotelo, P. A., 564
Souza, E., 1068
Spencer, T., 168
Spomer, N. A., 902
Stacy, E. A., 1068
Stark, J. D., 759
Stathas, G. J., 982
Steinkraus, D. C., 23
Stepić, S., 1204
Stevens, M., 1331
Stevenson, D. E., 1584
Stoeckli, S., 81, 1341
Storer, N. P., 168
Strange, J. P., 668
Streett, D. A., 631
Strohbehn, R., 555
Su, N.-Y., 592, 885
Subramanyam, B., 989, 998, 1022, 1719
Suckling, D. M., 720, 1295
Sullivan, B. T., 1821
Sumerford, D. V., 1406
Sumner, P. E., 631
Sun, J.-H., 1146
Swan, K. S., 1140
Szendrei, Z., 1039

T

Tabashnik, B. E., 504, 927
Tajima, R., 1167
Talebi, K., 1432
Talmich, E. E., 541
Tansey, J. A., 159
Tasei, J.-N., 1737
Taylor, P. B., 1140
Tebbets, J. S., 1515
Terán-Vargas, A. P., 168
Throne, J. E., 605, 616, 631, 1974
Tindall, K. V., 492
Tizado, E. J., 1792
Todd, J. H., 1944
Tolman, J. H., 365, 375
Tomberlin, J. K., 480
Toscano, N. C., 174
Toyoumi, A., 880
Triapitsyn, S. V., 1771
Tropp, J. M., 838
Tsunoda, K., 880

U

Ueda, K., 1568
Uesugi, R., 1704
Underwood, R. M., 256, 1057
Unruh, T., 1919

V

Vadivambal, R., 1011
Valdovinos-Nuñez, G. R., 132
Vancova, M., 1028
Van Duyn, J. W., 1950
vanEngelsdorp, D., 256
van Herk, W. G., 365, 375
Vargas, R. I., 759
Vargo, E. L., 1349
Velis, G., 1075
Vernon, R. S., 365, 375
Villa, J. D., 1756
Villet, M. H., 99
Vincent, C., 118, 584, 1838
Vitagliano, S., 444

W

Wagner, M. R., 944
Wakamura, S., 1568
Walker, N. R., 1624
Wallin, K. F., 1826
Walter, G. H., 17
Wang, D.-L., 360
Wang, J.-J., 1711
Wang, Q., 1771
Wang, Y. H., 515
Ward, K., 654
Ward, R., 654
Warriner, R. A., 902
Warrington, C. V., 1471
Watmough, J., 1575
Way, M. O., 216, 237, 1233
Weiland, A. A., 569
Weill, M., 484
Weintraub, P. G., 1337
Weston, P. A., 1466
White, N.D.G., 1011
Whitney, S., 728
Wilen, J., 915
Wille, B. D., 1228
Willett, L., 1919
Williams, D. J., 334
Williams, K. K., 1253
Williams, T., 42
Wilson, L. T., 216, 237, 1501
Winston, M. L., 784
Wu, J.-C., 384
Wu, K., 1805
Wu, Q.-J., 1888
Wu, Z., 1897
Wyckhuys, K.A.G., 1805

X

Xiao, L.-S., 1711
Xie, Q., 736
Xin, N., 623
Xu, Z., 933
Xue, F., 1805

Y

Yacoub, R., 1041
Yafuso, C. M., 504, 927
Yamamura, K., 1568
Yan, S., 1897
Yang, E. C., 1743
Yang, M.-M., 1152
Yao, H.-W., 1415
Ye, G.-Y., 1415
Yeh, W.-B., 1152
Yewhalaw, D., 50
Yin, J.-L., 384
Yoshimura, T., 880
Youngman, R. R., 859
Youssef, N. N., 1158, 1831
Yu, J.-F., 384

Z

Zalom, F. G., 750
Zalucki, M. P., 61
Zhang, G., 182
Zhang, Q., 182
Zhang, R., 736
Zhang, Y.-J., 1888
Zhang, Y.-Z., 1146
Zhang, Z.-Y., 360
Zhao, Q., 1897
Zhou, W. J., 515
Zhou, X.-M., 1888
Zhu, K. Y., 1974
Zhu, S., 1471
Zhu, Y. C., 515, 933
Zhuang, Y. L., 515

CUMULATIVE KEY WORD INDEX
Journal of Economic Entomology
Volume 101, 2008

A

abamectin,
  functional expression, 1888
  GABA receptor, 1888
  Plutella xylostella, 1888
  Xenopus oocyte, 1888
abscission,
  Anthonomus grandis, 68
  bolls, 68
  squares, 68
Acarapis woodi, 654, 668, 1756
  Apis mellifera, 654, 668, 1756
  genetic resistance, 1756
  honey bees, 1756
  Nosema spp., 668
  package honey bees, 668
  Russian bees, 654
  tracheal mites, 1756
  Varroa destructor, 654, 668
  varroa sensitive hygiene, 654
acaricide resistance,
  cross-resistance, 1704
  hitchhiking effect, 1704
  multiple resistance, 1704
  Tetranychus urticae, 1704
acephate,
  Acer pseudoplatanus, 822
  elm spanworm, 822
  Ennomos subsignaria, 822
Acer pseudoplatanus,
  acephate, 822
  elm spanworm, 822
  Ennomos subsignaria, 822
acetic acid,
  deformed wing virus, 256
  formic acid, 256
  fumigation, 256
  varroa mites, 256
acrinathrin,
  fitness, 499
  insecticide resistance, 499
  spinosad, 499
  Thysanoptera, 499
adult,
  cold storage, 1760
  demographics, 1760
  Conatocerus ashmeadi, 1760
  maternal and progeny fitness, 1760
adult colonization,
  pea, 1606
  pea leaf weevil, 1606
  reduced tillage, 1606
African bollworm,
  attract-and-kill, 99
  Chicorium intybus, 99
  chicory, 99
  Helicoverpa armigera, 99
Agrilus planipennis,
  ash, 1643
  canopy thinning, 1643
  emerald ash borer, 1643
  Fraxinus, 1643
Agriotes obscurus,
  fipronil, 375
  insecticide toxicity, 375
  survival time, 375
  wireworm, 375
Alcaligenes xylosoxidans,
  arthropod vector, 1519
  paratransgenesis, 1519
  Pierce's disease, 1519
  symbiotic control, 1519
alfalfa leafcutting bee, 660, 674, 1088
  Apoidea, 674
  bees, 660, 1088
  chalkbrood, 660, 1088
  disease control, 660
  fungicide toxicity, 1088
  insect pathology, 660, 1088
  Medicago, 674
  Megachilidae, 674
  pollen ball, 674
allele frequency,
  Bt resistance, 909
  Cry2Ab, 909
  dominance, 909
  F_2 screen, 909
Allium cepa,
  onion thrips, 1331
  Thrips tabaci, 1331
  within-plant distribution, 1331
almond leaf scorch,
  grass sharpshooter, 1105
  Pierce's disease (PD) of grapevines, 1105
  population dynamics, 1105
  transmission efficiency, 1105
almonds,
  Amyelois transitella, 1633
  mating disruption, 1633
  pistachios, 1633
Alphitobius diaperinus, 61, 480
  insect pest, 61
  insecticide resistance, 480
  knockdown, 480
  lesser mealworm, 480
  movement, 61
  poultry litter, 61
  recovery, 480
  Tenebrionidae, 61
American foulbrood disease,
  Apis mellifera, 1095
  Paenibacillus larvae, 1095
  shaking technique, 1095
  shook-swarm, 1095
Amorbia emigratella,
  phytosanitary treatment, 710
  postharvest, 710
  quarantine pest, 710
Amyelois transitella,
  almonds, 1633
  mating disruption, 1633
  pistachios, 1633
Anagasta kuehniella,
  biological control, 982
  Habrobracon hebetor, 982
  life tables, 982
  Plodia interpunctella, 982
Anarsia lineatella,
  bionomics, 1557
  developmental rate, 1557
  model evaluation, 1557
  temperature, 1557
Anastrepha ludens,
  commodity treatment, 716
  disinfestation, 716
  quarantine, 716
  radiation, 716
ant baits,
  ant management, 1068
  invasive species, 1068
  little fire ant, 1068
  unicolonial, 1068
ant management,
  ant baits, 1068
  invasive species, 1068
  little fire ant, 1068
  unicolonial, 1068
Anthonomus grandis,
  abscission, 68
  bolls, 68
  squares, 68
Anthonomus grandis grandis, 1600, 1729
  boll weevil, 1729
  dispersal, 1729
  eradication, 1729
  Gossypium hirsutum, 1600
  harvest, 1600
  insecticide, 1600
  population genetics, 1729

proactive, 1600
antiaggregation pheromones,
Dendroctonus pseudotsugae, 1826
3-methylcyclohex-2-en-1-one, 1826
antibiosis, 564, 1341
Aphis pomi, 1341
apple, 1341
Brachiaria, 564
quantitative trait loci, 1341
resistance, 564
shoot growth characteristics, 1341
spittlebugs, 564
sublethal effects, 564
Apanteles sp.,
Brachycyrtus sp., 692
life table, 692
Macrocentrus sp., 692
oviposition choice tests, 692
aphicide,
Leptinotarsa decemlineata, 74
Solanum tuberosum, 74
aphid predators,
conservation biocontrol, 1526
insectary crops, 1526
organic agriculture, 1526
Syrphidae, 1526
Aphis glycines, 278, 391, 801, 1228, 1488
artificial diet, 1228
Bean pod mottle virus, 801
Cerotoma trifurcata, 801
classical biological control, 278
coefficient of variation, 1488
Glycine max, 801
Harmonia axyridis, 391
insecticides, 391
nutrition, 1228
organic, 391
parasitoids, 278
plant phenology, 1488
predatory flies, 278
sampling, 1488
soybean, 391
soybean aphid, 1228
Aphis gossypii,
biological control, 23
epizootics, 23
Neozygites fresenii, 23
Aphis pomi,
antibiosis, 1341
apple, 1341
quantitative trait loci, 1341
shoot growth characteristics, 1341
Apis,
bee breeding, 272
feral honey bees, 272
population genetics, 272
Apis mellifera, 654, 668, 1057, 1075, 1081, 1095, 1756
Acarapis woodi, 654, 668, 1756
American foulbrood disease, 1095
desert conditions, 1057
drones, 1081
genetic resistance, 1756
honey bee, 1081, 1756
miticides, 1081
Nosema spp., 668
oxalic acid, 1057
package honey bees, 668
Paenibacillus larvae, 1095
PVC strips, 1075
residues, 1075
rotenone, 1075
Russian bees, 654
shaking technique, 1095
shook-swarm, 1095
spermatozoa, 1081
Sucrocide, 1057
tracheal mites, 1756
Varroa destructor, 654, 668, 1057, 1075
varroa sensitive hygiene, 654
Apoidea,
alfalfa leafcutting bee, 674
Medicago, 674
Megachilidae, 674
pollen ball, 674
Apolygus lucorum,
endosulfan, 1805
sublethal effects, 1805
toxicity, 1805
apple, 145, 206, 1341
antibiosis, 1341
Aphis pomi, 1341
beneficials, 145
control, 145
germplasm, 206
host plant resistance, 206
kaolin, 145
pests, 145
quantitative trait loci, 1341
Rhagoletis pomonella, 206
shoot growth characteristics, 1341
apple pests,
Cydia pomonella, 190
gene flow, 190
insecticide resistance, 190
microsatellites, 190
Argentine ant,
Linepithema humile, 699
liquid baits, 699
Pseudococcidae, 699
vineyards, 699
Argopistes tsekooni,
flea beetle, 1146
host specificity, 1146
Ligustrum sinense, 1146
weed biological control, 1146
armored scale insects,
cytochrome oxidase I and *II* genes, 1944
distribution and species composition, 1944
DNA diagnostic, 1944
multiplex PCR, 1944
arthropod vector,
Alcaligenes xylosoxidans, 1519
paratransgenesis, 1519
Pierce's disease, 1519
symbiotic control, 1519
artificial diet, 616, 1127, 1228
Aphis glycines, 1228
booklice, 616
fecundity, 1127
nutrition, 1228
Orius laevigatus, 1127
Psocoptera, 616
quality assurance, 1127
sex ratio, 616
soybean aphid, 1228
stored products, 616
ash,
Agrilus planipennis, 1643
canopy thinning, 1643
emerald ash borer, 1643
Fraxinus, 1643
assimilate allocation,
Nilaparvata lugens, 384
physiological response, 384
rice, 384
Atimia confusa confusa,
Callidiellum rufipenne, 430
Callidium frigidum, 430
Semanotus ligneus ligneus, 430
trap log, 430
atmospheric pressure plasma,
dielectric barrier discharge, 302
insect control, 302
oxygenated helium plasma, 302
quarantine, 302
attract-and-kill system, 99, 352
African bollworm, 99
Bactrocera dorsalis, 352
Chicorium intybus, 99
chicory, 99
Helicoverpa armigera, 99
methyl eugenol, 352
neonicotinoid insecticides, 352
oriental fruit fly, 352
attractant, 126, 769
Ceratitis capitata, 126
fruit fly, 126
(9*E*,11*Z*)-hexadecadien-1-yl acetate, 769
Lepidoptera, 769
lure, 126
pheromone antagonist, 769

pheromone race, 769
trap, 126
attractant synergist,
bark beetles, 1821
endo-brevicomin, 1821
monitoring, 1821
pheromone, 1821
attractiveness,
Oebalus pugnax, 1233
Oryza sativa, 1233
rice panicle stage, 1233
rice stink bug, 1233
augmentation biological control,
microclimate, 421
predation, 421
Rhyacionia frustrana, 421
Trichogramma exiguum, 421
autosomal,
esterases, 472
incompletely dominant, 472
indoxacarb, 472
Spodoptera litura, 472
avocado pests,
Braconidae, 1310
Ichneumonidae, 1310
Persea americana, 1310
quarantine, 1310

B

Bacillus thuringiensis, 36, 182, 1691, 1950
baseline susceptibility, 1691
benzylideneacetone, 36
bioassay, 1691
bollworm, 1950
Cnaphalocrocis medinalis, 1691
cotton, 1950
cry1Ac, 182
cry2A, 182
cry9C, 182
feeding behavior, 182
immunosuppression, 36
integrated pest management, 1950
microaggregation, 36
Spodoptera exigua, 36
tobacco budworm, 1950
Bacillus thuringiensis corn,
corn hybrids, 555
insect injury, 555
western bean cutworm, 555
Bactrocera cucurbitae,
Bactrocera dorsalis, 759
fruit fly suppression, 759
male annihilation, 759
SPLAT, 759
Bactrocera dorsalis, 352, 736, 759
attract-and-kill system, 352
Bactrocera cucurbitae, 759
eclosion time, 736
fruit fly suppression, 759
heat exposure, 736
male annihilation, 759
methyl eugenol, 352
neonicotinoid insecticides, 352
oriental fruit fly, 352
puparial morphology, 736
SPLAT, 759
timing of death, 736
baculovirus,
biopesticide, 1540
pesticide residue, 1540
PoGV, 1540
Solanum tuberosum, 1540
bait, 151, 873, 894
[^{14}C]hexaflumuron, 873
Blattella germanica, 151, 894
carbohydrates, 873
feeding, 873
German cockroach, 894
horizontal transfer, 894
indoxacarb, 894
sticky trap, 151
trap bias, 151
trapping, 151
xylose, 873
bark beetle, 829, 1821
attractant synergist, 1821
distribution, 829
endo-brevicomin, 1821
host range, 829
invasive species, 829
monitoring, 1821
pheromone, 1821
Pinus, 829
barley,
disinfestation, 1011
microwaves, 1011
mortality, 1011
stored-grain insects, 1011
baseline susceptibility,
Bacillus thuringiensis, 1691
bioassay, 1691
Cnaphalocrocis medinalis, 1691
Bean pod mottle virus,
Aphis glycines, 801
Cerotoma trifurcata, 801
Glycine max, 801
beat bucket,
beneficial arthropods, 1501
cotton, 1501
Orius, 1501
spiders, 1501
bee breeding,
Apis, 272
feral honey bees, 272
population genetics, 272
bees, 132, 660, 1088
alfalfa leafcutting bee, 660, 1088
chalkbrood, 660, 1088
crop pollination, 132
disease control, 660
fungicide toxicity, 1088
greenhouse, 132
insect pathology, 660, 1088
Yucatán, 132
behavior,
oviposition, 118
particle film technology, 118
Rhagoletis mendax, 118
Surround WP, 118
behavioral effects,
orchard pollination, 647
side effects, 647
sublethal dosages, 647
Bemisia argentifolii,
imidacloprid-resistant whiteflies, 174
Oberon, 174
Q Biotype, 174
whitefly immature stages, 174
Bemisia tabaci,
haplodiploid, 927
insect growth regulator, 927
insecticide resistance, 927
pyriproxyfen, 927
beneficial arthropods,
beat bucket, 1501
cotton, 1501
Orius, 1501
spiders, 1501
beneficials,
apples, 145
control, 145
kaolin, 145
pests, 145
Benin,
cassava chips, 1967
infestation, 1967
Prostephanus truncatus, 1967
yam chips, 1967
benzylideneacetone,
Bacillus thuringiensis, 36
immunosuppression, 36
microaggregation, 36
Spodoptera exigua, 36
bifenthrin,
fipronil, 1373
half-life, 1373
imidacloprid, 1373
toxicity, 1373
binomial data,
IPM, 1719
monitoring, 1719
sequential sampling, 1719
bioassay,
Bacillus thuringiensis, 1691
baseline susceptibility, 1691
Cnaphalocrocis medinalis, 1691
biofix,
codling moth, 1651
phenology, 1651
biological agent,
Liriomyza sativae, 1510
Liriomyza trifolii, 1510

multiplex polymerase chain reaction, 1510
Neochrysocharis formosa, 1510
biological control, 23, 251, 982, 1140, 1771
Anagasta kuehniella, 982
Aphis gossypii, 23
collection strategy, 1771
egg parasitoid, 1771
entomopathogenic nematodes, 251
epizootics, 23
eucalyptus longhorned borer, 1771
Habrobracon hebetor, 982
life tables, 982
microbial control, 251
Neozygites fresenii, 23
parasitic Hymenoptera, 1140
Phoracantha, 1771
Plodia interpunctella, 982
sex determination, 1140
sex ratio, 1140
strawberry, 251
bionomics,
Anarsia lineatella, 1557
developmental rate, 1557
model evaluation, 1557
temperature, 1557
biopesticide,
baculovirus, 1540
pesticide residue, 1540
PoGV, 1540
Solanum tuberosum, 1540
biosecurity,
internal transcribed spacer, 1167
nuclear ribosomal RNA gene, 1167
restriction fragment length polymorphism, 1167
biotechnology,
Bt cotton, 915
refuge, 915
refugia, 915
resistance, 915
biotype, 569, 955
Diuraphis noxia, 569
intrinsic rate of increase, 955
plant resistance, 569
Russian wheat aphid, 955
Triticum aesticum, 569
biting flies,
cloth targets, 1034
fly control, 1034
livestock pests, 1034
black grass bugs,
Capsus, 309
grazing management, 309
Irbisia, 309
Labops, 309
black widow spider,
controlled atmosphere, 1515
postharvest pest control, 1515
table grape, 1515
ultralow oxygen, 1515
Blattella germanica, 151, 894, 1384, 1389
bait, 151, 894
Camponotus floridanus, 1389
Cimex lectularius, 1389
dog, 1389
fatty acid salts, 1384
German cockroach, 894
horizontal transfer, 894
indoxacarb, 894
low toxic, 1384
Periplaneta americana, 1384
pseudoscent, 1389
soap, 1384
sticky trap, 151
trap bias, 151
trapping, 151
Blissus occiduus,
chinch bug, 533
plant resistance, 533
turfgrass, 533
bloom,
dispersal, 56
integrated pest management, 56
refuge, 56
boll weevil, 399, 1729
Anthonomus grandis grandis, 1729
dichlorvos, 399
dispersal, 1729
eradication, 1729
grandlure, 399
monitoring, 399
population genetics, 1729
trapping, 399
Bollgard,
Bollgard II, 546
Cry1Ac, 546
Cry2Ab2, 546
Lepidoptera, 546
Bollgard II, 546, 1594
Bollgard, 546
Bollworm, 1594
Cry1Ac, 546
Cry2Ab2, 546
Helicoverpa, 1594
integrated pest management, 1594
Lepidoptera, 546
Widestrike, 1594
bolls,
abscission, 68
Anthonomus grandis, 68
squares, 68
bollworm, 1244, 1594, 1950
Bacillus thuringiensis, 1950
Bollgard II, 1594
cotton, 1950
Helicoverpa, 1594
integrated pest management, 1594, 1950
tobacco budworm, 1244, 1950
transgenic cotton, 1244
Vip3A, 1244
VipCot, 1244
Widestrike, 1594
Bombus terrestris,
comparative bioassay, 1737
nutritive value, 1737
pollen consumption, 1737
pollen diet, 1737
booklice,
artificial diets, 616
Psocoptera, 616
sex ratio, 616
stored products, 616
botanical insecticide,
cinnamaldehyde, 1960
natural fumigant, 1960
Sitophilus oryzae, 1960
structure–activity relationship, 1960
Brachiaria,
antibiosis, 564
resistance, 564
spittlebugs, 564
sublethal effects, 564
Brachycyrtus sp.,
Apanteles sp., 692
life table, 692
Macrocentrus sp., 692
oviposition choice tests, 692
Braconidae,
avocado pests, 1310
Ichneumonidae, 1310
Persea americana, 1310
quarantine, 1310
Brassica campestris subsp. *napus* variety *pekinensis*,
Bt transgenic Chinese cabbage, 1134
enzyme-linked immunosorbent assay, 1134
Mamestra brassicae, 1134
Microplitis mediator, 1134
broccoli,
Frankliniella occidentalis, 1786
lettuce, 1786
phosphine, 1786
quarantine control, 1786
brood pheromone,
honey bee, 1749
winter colony growth, 1749

Bt corn, 492, 1614
cornstalk borers, 492
Diatraea saccharalis, 492
European corn borer, 1614
insecticide resistance management, 492
Ostrinia nubilalis, 1614
remote sensing, 1614
resistance, 1614
resistance monitoring, 492
Bt cotton, 504, 915, 933
biotechnology, 915
economic threshold, 959
F_1 screen, 933
Helicoverpa armigera, 933
host plant resistance, 959
Pectinophora gossypiella, 504
pink bollworm, 504
potato leafhopper, 959
refuge, 504, 915
refugia, 915
resistance, 504, 915
resistance detection, 933, 959
resistance management, 933
sap-feeding insect, 959
tolerance, 959
Bt resistance,
allele frequency, 909
Cry2Ab, 909
dominance, 909
F_2 screen, 909
Bt transgenic Chinese cabbage,
Brassica campestris subsp. *napus* variety *pekinensis*, 1134
enzyme-linked immunosorbent assay, 1134
Mamestra brassicae, 1134
Microplitis mediator, 1134
Buprestidae,
Cerambycidae, 409
Monochamus clamator, 409
semiochemicals, 409
Siricidae, 409
Busseola fusca,
maize, 90
nitrogen and potassium, 90
stemborer, 90
yield loss, 90

C

cabbage,
insecticide, 438
onion thrips, 438
plant resistance, 438
Thrips tabaci, 438
Cacopsylla chinensis,
cytochrome oxidase I, 1152
invasive species, 1152
16S rDNA, 1152
restriction fragment length polymorphism, 1152
cactus moth,
circadian rhythm, 314
dispersal, 314
flight, 314
sterile insect technique, 314
calcofluor,
chitinase, 1028
diflubenzuron, 1028
mites, 1028
peritrophic membrane, 1028
California red scale,
citricola scale, 451
citrus red mite, 451
cottony cushion scale, 451
neonicotinoid, 451
Callidiellum rufipenne,
Atimia confusa confusa, 430
Callidium frigidum, 430
Semanotus ligneus ligneus, 430
trap log, 430
Callidium frigidum,
Atimia confusa confusa, 430
Callidiellum rufipenne, 430
Semanotus ligneus ligneus, 430
trap log, 430
Camponotus floridanus,
Blattella germanica, 1389
Cimex lectularius, 1389
dog, 1389
pseudoscent, 1389
Canada thistle,
Cirsium arvense, 720
floral volatiles, 720
invasive species, 720
kairomone, 720
canola, 159, 810, 1811
cultural control, 810
decreased pesticide load, 1811
flea beetle injury, 1811
flea beetles, 159, 810
neonicotinoid, 159
pest management, 810
Phyllotreta spp., 810, 1811
seed treatment, 159, 1811
species composition shift, 159
canopy thinning,
Agrilus planipennis, 1643
ash, 1643
emerald ash borer, 1643
Fraxinus, 1643
Capsus,
black grass bugs, 309
grazing management, 309
Irbisia, 309
Labops, 309
carbamates,
organophosphates, 1667
pyrethroids, 1667
resistance, 1667
Spodoptera litura, 1667
carbohydrates,
bait, 873
feeding, 873
[^{14}C]hexaflumuron, 873
xylose, 873
Cardinium,
development, 1711
life table parameters, 1711
Liposcelis bostrychophila, 1711
reproduction, 1711
cassava chips,
Benin, 1967
infestation, 1967
Prostephanus truncatus, 1967
yam chips, 1967
CATTS,
codling moth, 295
controlled atmospheres, 295
heat, 295
oriental fruit moth, 295
Cerambycidae, 107, 409
Buprestidae, 409
Curculionidae, 107
detection, 107
Monochamus clamator, 409
multiple-funnel trap, 107
Scolytidae, 107
semiochemicals, 409
Siricidae, 409
Ceratitis capitata, 126, 866, 1325
attractant, 126
citrus peel, 866
development, 866
dispenser, 1325
fruit fly, 126, 1325
host plant resistance, 866
lure, 126
pheromone, 1325
survival, 866
trap, 126
trimedlure, 1325
cereals,
electrical penetration graph, 9
fecundity, 9
life table, 9
Myzus persicae, 9
Cerotoma trifurcata, 801, 1211
Aphis glycines, 801
Bean pod mottle virus, 801
Glycine max, 801, 1211
pest management, 1211
seed treatment, 1211
Chaetocnema confinis,
harvest date, 1238
planting date, 1238
soil temperature, 1238
sweetpotato, 1238
chalkbrood, 660, 1088
alfalfa leafcutting bee, 660, 1088
bees, 660, 1088
disease control, 660

fungicide toxicity, 1088
insect pathology, 660, 1088
Chicorium intybus,
African bollworm, 99
attract-and-kill, 99
chicory, 99
Helicoverpa armigera, 99
chicory,
African bollworm, 99
attract-and-kill, 99
Chicorium intybus, 99
Helicoverpa armigera, 99
Chilo partellus,
COI, 686
Cotesia flavipes, 686
Ethiopia, 686
sugarcane, 686
Chilo suppressalis,
fipronil, 1415
metabolism, 1415
Sesamia inferens, 1415
susceptibility, 1415
chinch bug,
Blissus occiduus, 533
plant resistance, 533
turfgrass, 533
chitinase,
calcofluor, 1028
diflubenzuron, 1028
mites, 1028
peritrophic membrane, 1028
Chrysoperla carnea,
organophosphates, 1676
pyrethroids, 1676
resistance, 1676
stability of resistance, 1676
Cicadellidae,
clean stock program, 1779
Diaspididae, 1779
quarantine, 1779
Tettigoniidae, 1779
Cicadellinae,
citrus variegated chlorosis, 1114
coffee leaf scorch, 1114
leafhopper vectors, 1114
Cimex lectularius,
Blattella germanica, 1389
Camponotus floridanus, 1389
dog, 1389
pseudoscent, 1389
cinnamaldehyde,
botanical insecticide, 1960
natural fumigant, 1960
Sitophilus oryzae, 1960
structure–activity relationship, 1960
circadian rhythm,
cactus moth, 314
dispersal, 314
flight, 314
sterile insect technique, 314
Circulifer spp.,
Exitianus capicola, 1337
Orosius orientalis, 1337
physical control, 1337
UV-absorbing plastic, 1337
Cirsium arvense,
Canada thistle, 720
floral volatiles, 720
invasive species, 720
kairomone, 720
citricola scale,
California red scale, 451
citrus red mite, 451
cottony cushion scale, 451
neonicotinoid, 451
citrus peel,
Ceratitis capitata, 866
development, 866
host plant resistance, 866
survival, 866
citrus red mite,
California red scale, 451
citricola scale, 451
cottony cushion scale, 451
neonicotinoid, 451
citrus variegated chlorosis,
Cicadellinae, 1114
coffee leaf scorch, 1114
leafhopper vectors, 1114
classical biological control,
Aphis glycines, 278
parasitoids, 278
predatory flies, 278
clean stock program,
Cicadellidae, 1779
Diaspididae, 1779
quarantine, 1779
Tettigoniidae, 1779
climatic conditions,
Dermolepida albohirtum, 334
pan evaporation, 334
sugarcane, 334
time series, 334
cloth targets,
biting flies, 1034
fly control, 1034
livestock pests, 1034
Cnaphalocrocis medinalis,
Bacillus thuringiensis, 1691
baseline susceptibility, 1691
bioassay, 1691
Coccinellidae,
Hippodamia variegata, 1792
marginal vegetation, 1792
supplemental food, 1792
Cochylis hospes,
economic distance, 969
EIL, 969
Helianthus annuus, 969
sampling, 969
codling moth, 295, 1651
biofix, 1651
CATTS, 295
controlled atmospheres, 295
heat, 295
oriental fruit moth, 295
phenology, 1651
coefficient of variation,
Aphis glycines, 1488
plant phenology, 1488
sampling, 1488
coffee leaf scorch,
Cicadellinae, 1114
citrus variegated chlorosis, 1114
leafhopper vectors, 1114
COI,
Chilo partellus, 686
Cotesia flavipes, 686
Ethiopia, 686
sugarcane, 686
cold storage,
adult, 1760
demographics, 1760
Gonatocerus ashmeadi, 1760
maternal and progeny fitness, 1760
collection strategy,
biological control, 1771
egg parasitoid, 1771
eucalyptus longhorned borer, 1771
Phoracantha, 1771
colonization behavior,
electronic penetration graph, 728
Myzus persicae, 728
probing behavior, 728
Rhopalosiphum padi, 728
commodity treatment, 716, 1051
Anastrepha ludens, 716
disinfestation, 716
Plodia interpunctella, 1051
quarantine, 716, 1051
radiation, 716, 1051
Sitotroga cerealella, 1051
common smut,
corn earworm, 1455
husk extension, 1455
maize weevil, 1455
stink bugs, 1455
community ecology,
feed mills, 998
insect survey, 998
stored-product insects, 998
trapping, 998
comparative bioassay,
Bombus terrestris, 1737
nutritive value, 1737
pollen consumption, 1737
pollen diet, 1737

Conotrachelus nenuphar, 1302, 1838
integrated pest management, 1302
material effect, 1838
monitoring, 1838
pheromone, 1302
plum curculio, 1302
traps, 1838
weather effect, 1838
weevil, 1302
conservation biocontrol,
aphid predators, 1526
insectary crops, 1526
organic agriculture, 1526
Syrphidae, 1526
conservation tillage,
Ipomoea batatas, 1871
no-till, 1871
pitfall traps, 1871
control,
apples, 145
beneficials, 145
kaolin, 145
pests, 145
controlled atmospheres, 295, 1515
black widow spider, 1515
CATTS, 295
codling moth, 295
heat, 295
oriental fruit moth, 295
postharvest pest control, 1515
table grape, 1515
ultralow oxygen, 1515
Coptotermes formosanus, 592, 880, 1367
dwellings, 1367
feeding preference, 880
flooding, 1367
gamma-irradiated wood, 880
hexaflumuron, 880
Isoptera, 592
mark–recapture protocol, 592
noviflumuron, 880
population estimate, 592
probability density function, 592
termite, 1367
corn earworm, 1455, 1471
common smut, 1455
Glycine max, 1471
husk extension, 1455
maize weevil, 1455
marker-assisted selection, 1471
quantitative trait loci, 1471
soybean looper, 1471
stink bugs, 1455
corn hybrids,
Bacillus thuringiensis corn, 555
insect injury, 555
western bean cutworm, 555
corn pest,
monitoring, 404
pest management, 404
Striacosta albicosta, 404
corn rootworm,
physiological time model, 1584
cornstalk borers,
Bt corn, 492
Diatraea saccharalis, 492
insecticide resistance management, 492
resistance monitoring, 492
Cotesia flavipes, 50, 686
Chilo partellus, 686
COI, 686
Ethiopia, 686
maize, 50
stemborer, 50
sugarcane, 686
trap plants, 50
wild hosts, 50
cotton, 1501, 1950
Bacillus thuringiensis, 1950
beat bucket, 1501
beneficial arthropods, 1501
bollworm, 1950
integrated pest management, 1950
Orius, 1501
spiders, 1501
tobacco budworm, 1950
cottony cushion scale,
California red scale, 451
citricola scale, 451
citrus red mite, 451
neonicotinoid, 451
crop pollination,
bees, 132
greenhouse, 132
Yucatán, 132
cropping-structure,
insecticide-resistance, 1685
Thrips, 1685
cross-resistance, 1704, 1911
acaricide resistance, 1704
Cry1Ac, 1911
diamondback moth, 1911
genetics of resistance, 1911
hitchhiking effect, 1704
multiple resistance, 1704
pyrethroids, 1911
Tetranychus urticae, 1704
Cry1Ac, 182, 546, 1406, 1421, 1911
Bacillus thuringiensis, 182
Bollgard, 546
Bollgard II, 546
cross-resistance, 1911
cry2A, 182
Cry2Ab2, 546
cry9C, 182
diamondback moth, 1911
feeding behavior, 182
fitness costs, 1421
genetics of resistance, 1911
Helicoverpa zea, 1421
Heliothis virescens, 1406
heritability of resistance, 1421
insecticide resistance management, 1406
Lepidoptera, 546
pyrethroids, 1911
single-pair family, 1406
stability of resistance, 1421
YHD2, 1406
Cry2Ab, 182, 909
allele frequency, 909
Bacillus thuringiensis, 182
Bt resistance, 909
cry1Ac, 182
cry9C, 182
dominance, 909
F_2 screen, 909
feeding behavior, 182
Cry2Ab2,
Bollgard, 546
Bollgard II, 546
Cry1Ac, 546
Lepidoptera, 546
cry9C,
Bacillus thuringiensis, 182
cry1Ac, 182
cry2A, 182
feeding behavior, 182
Cryptorhynchus lapathi,
host plant resistance, 199
hybrid poplars, 199
Populus maximowiczii, 199
Culex,
insecticide, 1204
instar, 1204
mosquito, 1204
Ochlerotatus, 1204
Culex pipiens,
esterase, 484
insecticide resistance genes, 484
mosquito, 484
cultural control,
canola, 810
flea beetles, 810
pest management, 810
Phyllotreta spp., 810
cultural management,
habitat management, 1317
onions, 1317
straw mulch, 1317
Thrips tabaci, 1317
Cuphea,
Diabrotica virgifera, 797
western corn rootworm, 797
Curculionidae,
Cerambycidae, 107
detection, 107
multiple-funnel trap, 107

Scolytidae, 107
Cydia pomonella, 81, 190, 288
apple pests, 190
fruit phenology, 81
gene flow, 190
granulovirus, 288
insecticide resistance, 190
larval infestation, 81
microsatellites, 190
monitoring, 81
shelf life, 288
storage, 288
temperature, 288
within-canopy distribution, 81
Cylas formicarius elegantulus,
gamma radiation, 1198
mating behavior, 1198
sterile insect technique (SIT), 1198
sweetpotato weevil, 1198
Cynodon,
cytochrome c oxidase I, 1624
Phyllophaga, 1624
scarab flight periods, 1624
white grub, 1624
cytochrome c oxidase I,
Cynodon, 1624
Phyllophaga, 1624
scarab flight periods, 1624
white grub, 1624
cytochrome oxidase I,
Cacopsylla chinensis, 1152
invasive species, 1152
16S rDNA, 1152
restriction fragment length polymorphism, 1152
cytochrome oxidase I and *II* genes,
armored scale insects, 1944
distribution and species composition, 1944
DNA diagnostic, 1944
multiplex PCR, 1944

D

Daktulosphaira vitifoliae,
DNA, 1934
grapevine, 1934
molecular diagnostic test, 1934
pest detection, 1934
soil, 1934
damage,
germination rate, 114
mango pulp, 114
mango seeds, 114
mango weevil, 114
decreased pesticide load,
canola, 1811
flea beetle injury, 1811
Phyllotreta spp., 1811
seed treatments, 1811
deficit irrigation,
grape, 1287
leafhoppers, 1287
vineyard, 1287
water, 1287
defoliation,
elm leaf beetle, 944
leaf toughness, 944
trichomes, 944
Ulmus spp., 944
deformed wing virus,
acetic acid, 256
formic acid, 256
fumigation, 256
varroa mites, 256
Delphacidae,
Reoviridae, 17
Saccharum, 17
sugarcane, 17
deltamethrin,
fitness costs, 1927
Heliothis virescens, 1927
Indoxacarb, 1927
resistance, 1927
demographics,
adult, 1760
cold storage, 1760
Gonatocerus ashmeadi, 1760
maternal and progeny fitness, 1760
Dendroctonus,
Ips, 1253
multiple regression models, 1253
Pinus ponderosa, 1253
stand characteristics, 1253
Dendroctonus pseudotsugae,
antiaggregation pheromones, 1826
3-methylcyclohex-2-en-1-one, 1826
density thresholds,
Dicyphinae, 1864
integrated pest management, 1864
omnivory, 1864
plant tolerance, 1864
Dermolepida albohirtum,
climatic conditions, 334
pan evaporation, 334
sugarcane, 334
time series, 334
desert conditions,
Apis mellifera, 1057
oxalic acid, 1057
Sucrocide, 1057
Varroa destructor, 1057
detection,
Cerambycidae, 107
Curculionidae, 107
multiple-funnel trap, 107
Scolytidae, 107
development, 325, 541, 866, 1711, 1798
Cardinium, 1711
Ceratitis capitata, 866
citrus peel, 866
development thresholds, 1798
fecundity, 325
host plant resistance, 866
life table, 325
life table parameters, 1711
Liposcelis bostrychophila, 1711
reproduction, 541, 1711, 1798
Russian wheat aphid, 541
Scymnus subvillosus, 325
survival, 866, 1798
temperature, 541
thermal constants, 1798
development thresholds,
development, 1798
reproduction, 1798
survival, 1798
thermal constants, 1798
developmental rates, 605, 1557
Anarsia lineatella, 1557
bionomics, 1557
life history, 605
marking, 605
model evaluation, 1557
psocids, 605
stored products, 605
temperature, 1557
Diabrotica virgifera,
Cuphea, 797
western corn rootworm, 797
Diabrotica virgifera virgifera, 341, 1547
fatty acid, 341
feeding stimulants, 341
insect behavior, 341
nutritional ecology, 1547
plant–insect interactions, 1547
Zea mays, 341
diamondback moth,
cross-resistance, 1911
Cry1Ac, 1911
genetics of resistance, 1911
pyrethroids, 1911
Diaphorina citri,
dispersion, 1478
sample size, 1478
sampling units, 1478
Diaprepes abbreviatus L.,
genetic diversity, 1443
invasive species, 1443
mitochondrial markers, 1443
quarantine pest, 1443
Diaspididae,
Cicadellidae, 1779
clean stock program, 1779
quarantine, 1779
Tettigoniidae, 1779

Diatraea saccharalis, 492, 1903
Bt corn, 492
cornstalk borers, 492
insecticide resistance management, 492
resistance, 1903
resistance monitoring, 492
selection, 1903
sugarcane borer, 1903
tebufenozide, 1903
dichlorvos,
boll weevil, 399
grandlure, 399
monitoring, 399
trapping, 399
Dicyphinae,
density thresholds, 1864
integrated pest management, 1864
omnivory, 1864
plant tolerance, 1864
dielectric barrier discharge,
atmospheric pressure plasma, 302
insect control, 302
oxygenated helium plasma, 302
quarantine, 302
diflubenzuron,
calcofluor, 1028
chitinase, 1028
mites, 1028
peritrophic membrane, 1028
disease control,
alfalfa leafcutting bee, 660
bees, 660
chalkbrood, 660
insect pathology, 660
disease management,
epidemiology, 1
insect vector, 1
modeling, 1
disinfestation, 716, 1011
Anastrepha ludens, 716
barley, 1011
commodity treatment, 716
microwaves, 1011
mortality, 1011
quarantine, 716
radiation, 716
stored-grain insects, 1011
dispenser,
Ceratitis capitata, 1325
fruit fly, 1325
pheromone, 1325
trimedlure, 1325
dispersal, 56, 314, 1729
Anthonomus grandis grandis, 1729
bloom, 56
boll weevil, 1729
cactus moth, 314
circadian rhythm, 314
eradication, 1729
flight, 314
integrated pest management, 56
population genetics, 1729
refuge, 56
sterile insect technique, 314
dispersion,
Diaphorina citri, 1478
sample size, 1478
sampling units, 1478
distribution,
bark beetle, 829
host range, 829
invasive species, 829
Pinus, 829
distribution and species composition,
armored scale insects, 1944
cytochrome oxidase I and *II* genes, 1944
DNA diagnostic, 1944
multiplex PCR, 1944
Diuraphis noxia,
biotype, 569
plant resistance, 569
Triticum aestivum, 569
DNA,
Daktulosphaira vitifoliae, 1934
grapevine, 1934
molecular diagnostic test, 1934
pest detection, 1934
soil, 1934
DNA barcoding,
DNA extraction, 523
invasive species, 523
quarantine pest, 523
species identification, 523
DNA diagnostic,
armored scale insects, 1944
cytochrome oxidase I and *II* genes, 1944
distribution and species composition, 1944
multiplex PCR, 1944
DNA extraction,
DNA barcoding, 523
invasive species, 523
quarantine pest, 523
species identification, 523
(*E,Z*)-7,9-dodecadienyl acetate,
(*E,Z*)-7,9-dodecadienyl acetate, 444
false-trail following, 444
grapevine, 444
mating disruption, 444
dog,
Blattella germanica, 1389
Camponotus floridanus, 1389
Cimex lectularius, 1389
pseudoscent, 1389
dominance,
allele frequency, 909
Bt resistance, 909
Cry2Ab, 909
F_2 screen, 909
drones,
Apis mellifera, 1081
honey bee, 1081
miticides, 1081
spermatozoa, 1081
dwellings,
Coptotermes formosanus, 1367
flooding, 1367
termite, 1367

E

eclosion time,
Bactrocera dorsalis, 736
heat exposure, 736
puparial morphology, 736
timing of death, 736
eco-races,
genetic variability, 1176
morphological traits, 1176
polymorphism, 1176
tasar silk worm, 1176
economic distance,
Cochylis hospes, 969
EIL, 969
Helianthus annuus, 969
sampling, 969
economic threshold,
Bt cotton, 959
F_1 screen, 959
host plant resistance, 959
potato leafhopper, 959
resistance detection, 959
sap-feeding insect, 959
tolerance, 959
egg parasitoid,
biological control, 1771
collection strategy, 1771
eucalyptus longhorned borer, 1771
Phoracantha, 1771
eggplant,
Leptinotarsa decemlineata, 859
potato, 859
solanaceous, 859
tomato, 859
EIL,
Cochylis hospes, 969
economic distance, 969
Helianthus annuus, 969
sampling, 969
Elateridae,
insecticide toxicity, 365
morbidity, 365
neonicotinoids, 365
wireworms, 365

electrical penetration graph, 9, 584, 728, 1183
cereals, 9
colonization behavior, 728
fecundity, 9
feeding behavior, 584
Homalodisca coagulata, 1183
host plant acceptance, 1183
life table, 9
Myzus persicae, 9, 728
olfactometry, 584
probing, 1183
probing behavior, 728
Rhopalosiphum padi, 728
Solanum chomatophilum, 584
Solanum stoloniferum, 584
elm leaf beetle,
defoliation, 944
leaf toughness, 944
trichomes, 944
Ulmus spp., 944
elm spanworm,
acephate, 822
Acer pseudoplatanus, 822
Ennomos subsignaria, 822
emerald ash borer, 1643, 1831
Agrilus planipennis, 1643
ash, 1643
canopy thinning, 1643
Fraxinus, 1643
prism trap, 1831
purple, 1831
trap height, 1831
visual cues, 1831
emergence,
landscape, 1575
Leptinotarsa decemlineata, 1575
sampling, 1575
endo-brevicomin,
attractant synergist, 1821
bark beetles, 1821
monitoring, 1821
pheromone, 1821
endosulfan,
Apolygus lucorum, 1805
sublethal effects, 1805
toxicity, 1805
Ennomos subsignaria,
acephate, 822
Acer pseudoplatanus, 822
elm spanworm, 822
entomopathogenic fungus,
epizootic, 885
Metarhizium, 885
termite, 885
entomopathogenic nematodes,
biological control, 251
microbial control, 251
strawberry, 251
enzyme-linked immunosorbent assay,
Brassica campestris subsp. *napus* variety *pekinensis*, 1134
Bt transgenic Chinese cabbage, 1134
Manestra brassicae, 1134
Microplitis mediator, 1134
epidemiology, 1, 1041
disease management, 1
Homalodisca vitripennis, 1041
insect vector, 1
modeling, 1
Pierce's disease, 1041
Xylella fastidiosa, 1041
epizootic, 23, 885
Aphis gossypii, 23
biological control, 23
entomopathogenic fungus, 885
Metarhizium, 885
Neozygites fresenii, 23
termite, 885
eradication, 838, 1729
Anthonomus grandis grandis, 1729
boll weevil, 1729
dispersal, 1729
invasive species, 838
population genetics, 1729
quarantine, 838
esterase, 472, 484
autosomal, 472
Culex pipiens, 484
incompletely dominant, 472
indoxacarb, 472
insecticide resistance genes, 484
mosquito, 484
Spodoptera litura, 472
esterase alleles,
insecticide resistance, 1897
mosquitoes, 1897
Ethiopia,
Chilo partellus, 686
COI, 686
Cotesia flavipes, 686
sugarcane, 686
ethyl formate,
fumigation, 623
insect, 623
methyl isothiocyanate, 623
residue, 623
ethylene glycol,
parasitoid, 631
Plodia interpunctella, 631
Sitophilus spp., 631
trapping, 631
eucalyptus longhorned borer,
biological control, 1771
collection strategy, 1771
egg parasitoid, 1771
Phoracantha, 1771
European corn borer,
Bt corn, 1614
Ostrinia nubilalis, 1614
remote sensing, 1614
resistance, 1614
Exitianus capicola,
Circulifer spp., 1337
Orosius orientalis, 1337
physical control, 1337
UV-absorbing plastic, 1337
exotic species,
invasive species, 1276
Lauraceae, 1276
laurel wilt, 1276

F

F_1 screen, 933, 959
Bt cotton, 933, 959
economic threshold, 959
Helicoverpa armigera, 933
host plant resistance, 959
potato leafhopper, 959
resistance detection, 933, 959
resistance management, 933
sap-feeding insect, 959
tolerance, 959
F_2 screen,
allele frequency, 909
Bt resistance, 909
Cry2Ab, 909
dominance, 909
fall armyworm,
haplotype, 742
migration, 742
Spodoptera frugiperda, 742
false-trail following,
(*E,Z*)-7,9-dodecadienyl acetate, 444
grapevine, 444
mating disruption, 444
fatty acid,
Diabrotica virgifera virgifera, 341
feeding stimulants, 341
insect behavior, 341
Zea mays, 341
fatty acid salts,
Blattella germanica, 1384
low toxic, 1384
Periplaneta americana, 1384
soap, 1384
fecundity, 9, 325, 777, 1127
artificial diet, 1127
cereals, 9
development, 325
electrical penetration graph, 9
Galleria mellonella, 777
life table, 9, 325
lysozyme, 777
Myzus persicae, 9
Orius laevigatus, 1127

quality assurance, 1127
Scymnus subvillosus, 325
sodium tetraborate, 777
survival, 777
feed mills,
community ecology, 998
insect survey, 998
stored-product insects, 998
trapping, 998
feeding,
bait, 873
carbohydrates, 873
[^{14}C]hexaflumuron, 873
xylose, 873
feeding behavior, 182, 584
Bacillus thuringiensis, 182
cry1Ac, 182
cry2A, 182
cry9C, 182
electrical penetration graph, 584
olfactometry, 584
Solanum chomatophilum, 584
Solanum stoloniferum, 584
feeding preference,
Coptotermes formosanus, 880
gamma-irradiated wood, 880
hexaflumuron, 880
noviflumuron, 880
feeding stimulants,
Diabrotica virgifera virgifera, 341
fatty acid, 341
insect behavior, 341
Zea mays, 341
feral honey bees,
Apis, 272
bee breeding, 272
population genetics, 272
fertility life table,
fitness cost, 1432
Myzus persicae, 1432
resistance to insecticide, 1432
fipronil, 375, 1373, 1397, 1415
Agriotes obscurus, 375
bifenthrin, 1373
Chilo suppressalis, 1415
half-life, 1373
imidacloprid, 1373
insecticide toxicity, 375
Linepithema humile, 1397
metabolism, 1415
necrophoresis, 1397
Sesamia inferens, 1415
spot treatment, 1397
survival time, 375
susceptibility, 1415
toxicity, 1373
wireworm, 375
fitness,
acrinathrin, 499
insecticide resistance, 499
spinosad, 499
Thysanoptera, 499
fitness costs, 1421, 1927
Cry1Ac, 1421
deltamethrin, 1927
fertility life table, 1432
Helicoverpa zea, 1421
Heliothis virescens, 1927
heritability of resistance, 1421
Indoxacarb, 1927
Myzus persicae, 1432
resistance, 1927
resistance to insecticide, 1432
stability of resistance, 1421
flea beetle injury,
canola, 1811
decreased pesticide load, 1811
Phyllotreta spp., 1811
seed treatments, 1811
flea beetles, 159, 810, 1146
Argopistes tsekooni, 1146
canola, 159, 810
cultural control, 810
host specificity, 1146
Ligustrum sinense, 1146
neonicotinoid, 159
pest management, 810
Phyllotreta spp., 810
seed treatment, 159
species composition shift, 159
weed biological control, 1146
flight,
cactus moth, 314
circadian rhythm, 314
dispersal, 314
sterile insect technique, 314
flooding,
Coptotermes formosanus, 1367
dwellings, 1367
termite, 1367
floral volatiles,
Canada thistle, 720
Cirsium arvense, 720
invasive species, 720
kairomone, 720
fly control,
biting flies, 1034
cloth targets, 1034
livestock pests, 1034
foraging behavior,
honey bee, 1743
imidacloprid, 1743
sublethal dosage, 1743
formic acid,
acetic acid, 256
deformed wing virus, 256
fumigation, 256
varroa mites, 256
Frankliniella occidentalis,
broccoli, 1786
lettuce, 1786
phosphine, 1786
quarantine control, 1786
Fraxinus,
Agrilus planipennis, 1643
ash, 1643
canopy thinning, 1643
emerald ash borer, 1643
fruit fly, 126, 1325
attractant, 126
Ceratitis capitata, 126, 1325
dispenser, 1325
lure, 126
pheromone, 1325
trap, 126
trimedlure, 1325
fruit fly suppression,
Bactrocera cucurbitae, 759
Bactrocera dorsalis, 759
male annihilation, 759
SPLAT, 759
fruit phenology,
Cydia pomonella, 81
larval infestation, 81
monitoring, 81
within-canopy distribution, 81
fumigant,
Lycoriella ingenua, 139
plant essential oils, 139
fumigation, 256, 623
acetic acid, 256
deformed wing virus, 256
ethyl formate, 623
formic acid, 256
insect, 623
methyl isothiocyanate, 623
residue, 623
varroa mites, 256
functional expression,
abamectin, 1888
GABA receptor, 1888
Plutella xylostella, 1888
Xenopus oocyte, 1888
fungicide toxicity,
alfalfa leafcutting bee, 1088
bees, 1088
chalkbrood, 1088
insect pathology, 1088

G

GABA receptor,
abamectin, 1888
functional expression, 1888
Plutella xylostella, 1888
Xenopus oocyte, 1888
Galleria mellonella,
fecundity, 777
lysozyme, 777
sodium tetraborate, 777
survival, 777
gamma radiation,
Cylas formicarius elegantulus, 1198
mating behavior, 1198

sterile insect technique (SIT), 1198
sweetpotato weevil, 1198
gamma-irradiated wood,
Coptotermes formosanus, 880
feeding preference, 880
hexaflumuron, 880
noviflumuron, 880
gene flow,
apple pests, 190
Cydia pomonella, 190
insecticide resistance, 190
microsatellites, 190
genetic diversity,
Diaprepes abbreviatus L., 1443
invasive species, 1443
mitochondrial markers, 1443
quarantine pest, 1443
genetic resistance,
Acarapis woodi, 1756
Apis mellifera, 1756
honey bees, 1756
tracheal mites, 1756
genetic variability,
eco-races, 1176
morphological traits, 1176
polymorphism, 1176
tasar silk worm, 1176
genetics,
Nilaparvata lugens, 575
resistance, 575
rice mutants, 575
genetics of resistance,
cross-resistance, 1911
Cry1Ac, 1911
diamondback moth, 1911
pyrethroids, 1911
genotyping,
microsatellite molecular markers, 1062
plowing, 1062
Solenopsis invicta, 1062
German cockroach,
bait, 894
Blattella germanica, 894
horizontal transfer, 894
indoxacarb, 894
germination rate,
damage, 114
mango pulp, 114
mango seeds, 114
mango weevil, 114
germplasm,
apple, 206
host plant resistance, 206
Rhagoletis pomonella, 206
glassy-winged sharpshooter,
orange trees, 226
pest management, 226
sex ratio, 226
yellow sticky trap, 226
Glycine max, 801, 1211, 1471
Aphis glycines, 801
Bean pod mottle virus, 801
Cerotoma trifurcata, 801, 1211
corn earworm, 1471
marker-assisted selection, 1471
pest management, 1211
quantitative trait loci, 1471
seed treatment, 1211
soybean looper, 1471
Gonatocerus ashmeadi,
adult, 1760
cold storage, 1760
demographics, 1760
maternal and progeny fitness, 1760
Gossypium hirsutum,
Anthonomus grandis grandis, 1600
harvest, 1600
insecticide, 1600
proactive, 1600
grains,
Rhyzopertha dominica, 976
Sitophilus oryzae, 976
spinosad, 976
grandlure,
boll weevil, 399
dichlorvos, 399
monitoring, 399
trapping, 399
granule formulation,
mating disruption, 1856
oriental beetle, 1856
soywax, 1856
turfgrass, 1856
granulovirus,
Cydia pomonella, 288
shelf life, 288
storage, 288
temperature, 288
grape,
deficit irrigation, 1287
leafhoppers, 1287
vineyard, 1287
water, 1287
grapevine, 444, 1934
Daktulosphaira vitifoliae, 1934
DNA, 1934
(*E,Z*)-7,9-dodecadienyl acetate, 444
false-trail following, 444
mating disruption, 444
molecular diagnostic test, 1934
pest detection, 1934
soil, 1934
grass sharpshooter,
almond leaf scorch, 1105
Pierce's disease (PD) of grapevines, 1105
population dynamics, 1105
transmission efficiency, 1105
grazing management,
black grass bugs, 309
Capsus, 309
Irbisia, 309
Labops, 309
greenhouse,
bees, 132
crop pollination, 132
Yucatán, 132

H

habitat management,
cultural management, 1317
onions, 1317
straw mulch, 1317
Thrips tabaci, 1317
Habrobracon hebetor,
Anagasta kuehniella, 982
biological control, 982
life tables, 982
Plodia interpunctella, 982
half-life,
bifenthrin, 1373
fipronil, 1373
imidacloprid, 1373
toxicity, 1373
haplodiploid,
Bemisia tabaci, 927
insect growth regulator, 927
insecticide resistance, 927
pyriproxyfen, 927
haplotype,
fall armyworm, 742
migration, 742
Spodoptera frugiperda, 742
Harmonia axyridis,
Aphis glycines, 391
insecticides, 391
organic, 391
soybean, 391
harvest,
Anthonomus grandis grandis, 1600
Gossypium hirsutum, 1600
insecticide, 1600
proactive, 1600
harvest date,
Chaetocnema confinis, 1238
planting date, 1238
soil temperature, 1238
sweetpotato, 1238
heat,
CATTS, 295
codling moth, 295
controlled atmospheres, 295
oriental fruit moth, 295
heat exposure,
Bactrocera dorsalis, 736
eclosion time, 736
puparial morphology, 736
timing of death, 736

heat shock proteins,
heat stress, 1974
heat treatment, 1974
Lepinotus reticulatus, 1974
Liposcelis entomophila, 1974
heat stress,
heat shock proteins, 1974
heat treatment, 1974
Lepinotus reticulatus, 1974
Liposcelis entomophila, 1974
heat treatment, 989, 1974
heat shock proteins, 1974
heat stress, 1974
Lepinotus reticulatus, 1974
Liposcelis entomophila, 1974
modeling, 989
survival, 989
Tribolium confusum, 989
Helianthus annuus,
Cochylis hospes, 969
economic distance, 969
EIL, 969
sampling, 969
Helicoverpa,
Bollgard II, 1594
Bollworm, 1594
integrated pest management, 1594
Widestrike, 1594
Helicoverpa armigera, 99, 933
African bollworm, 99
attract-and-kill, 99
Bt cotton, 933
Chicorium intybus, 99
chicory, 99
F_1 screen, 933
resistance detection, 933
resistance management, 933
Helicoverpa zea,
Cry1Ac, 1421
fitness costs, 1421
heritability of resistance, 1421
stability of resistance, 1421
Heliothis virescens, 168, 1406, 1927
Cry1Ac, 1406
deltamethrin, 1927
fitness costs, 1927
Indoxacarb, 1927
insecticide, 168
insecticide resistance management, 1406
resistance, 1927
resistance management, 168
single-pair family, 1406
transgenic cotton, 168
YHD2, 1406
Hemiptera,
Pierce's disease, 1122
quantification, 1122
real-time polymerase chain reaction, 1122
transmission, 1122
heritability of resistance,
Cry1Ac, 1421
fitness costs, 1421
Helicoverpa zea, 1421
stability of resistance, 1421
Heterorhabditis bacteriophora,
multiple species approach, 1533
Otiorhynchus ligustici, 1533
Steinernema carpocapsae, 1533
Steinernema feltiae, 1533
(9*E*:11*Z*)-hexadecadien-1-yl acetate,
attractant, 769
(9*E*:11*Z*)-hexadecadien-1-yl acetate, 769
Lepidoptera, 769
pheromone antagonist, 769
pheromone race, 769
hexaflumuron,
Coptotermes formosanus, 880
feeding preference, 880
gamma-irradiated wood, 880
noviflumuron, 880
[^{14}C]hexaflumuron,
[^{14}C]hexaflumuron, 873
bait, 873
carbohydrates, 873
feeding, 873
xylose, 873
Hippodamia variegata,
Coccinellidae, 1792
marginal vegetation, 1792
supplemental food, 1792
hitchhiking effect,
acaricide resistance, 1704
cross-resistance, 1704
multiple resistance, 1704
Tetranychus urticae, 1704
Homalodisca coagulata,
electrical penetration graph (EPG), 1183
host plant acceptance, 1183
probing, 1183
Homalodisca vitripennis,
epidemiology, 1041
Pierce's disease, 1041
Xylella fastidiosa, 1041
honey bee, 1081, 1743, 1749, 1756
Acarapis woodi, 1756
Apis mellifera, 1081, 1756
brood pheromone, 1749
drones, 1081
foraging behavior, 1743
genetic resistance, 1756
imidacloprid, 1743
miticides, 1081
spermatozoa, 1081
sublethal dosage, 1743
tracheal mites, 1756
winter colony growth, 1749
horizontal transfer, 894 902
bait, 894
Blattella germanica, 894
German cockroach, 894
indoxacarb, 894
Reticulitermes flavipes, 902
termiticide, 902
host plant acceptance,
electrical penetration graph (EPG), 1183
Homalodisca coagulata, 1183
probing, 1183
host plant resistance, 199, 206, 866, 959, 1466
apple, 206
Bt cotton, 959
Ceratitis capitata, 866
citrus peel, 866
Cryptorhynchus lapathi, 199
development, 866
economic threshold, 959
F_1 screen, 959
germplasm, 206
hybrid poplars, 199
induced defense, 1466
induced resistance, 1466
Populus maximowiczii, 199
potato leafhopper, 959
resistance detection, 959
Rhagoletis pomonella, 206
sap-feeding insect, 959
survival, 866
tolerance, 959
host range,
bark beetle, 829
distribution, 829
invasive species, 829
Pinus, 829
host specificity,
Argopistes tsekooni, 1146
flea beetle, 1146
Ligustrum sinense, 1146
weed biological control, 1146
humidity,
Menacanthus stramineus, 637
Ornithonyssus sylviarum, 637
survival, 637
temperature, 637
husk extension,
common smut, 1455
corn earworm, 1455
maize weevil, 1455
stink bugs, 1455
hybrid,
imported fire ants, 265
repellent, 265
Solenopsis invicta Buren, 265
Solenopsis richteri, 265
hybrid poplars,
Cryptorhynchus lapathi, 199
host plant resistance, 199
Populus maximowiczii, 199

I
Ichneumonidae,
avocado pests, 1310
Braconidae, 1310
Persea americana, 1310
quarantine, 1310
imidacloprid, 515, 1373, 1743
bifenthrin, 1373
fipronil, 1373
foraging behavior, 1743
half-life, 1373
honey bee, 1743
Nilaparvata lugens, 515
resistance, 515
risk assessment, 515
sublethal dosage, 1743
susceptibilities, 515
toxicity, 1373
imidacloprid-resistant whiteflies,
Bemisia argentifolii, 174
Oberon, 174
Q Biotype, 174
whitefly immature stages, 174
immunosuppression,
Bacillus thuringiensis, 36
benzylideneacetone, 36
microaggregation, 36
Spodoptera exigua, 36
imported fire ants,
hybrid, 265
repellent, 265
Solenopsis invicta Buren, 265
Solenopsis richteri, 265
incompletely dominant,
autosomal, 472
esterases, 472
indoxacarb, 472
Spodoptera litura, 472
indoxacarb, 461, 472, 894, 1927
autosomal, 472
bait, 894
Blattella germanica, 894
deltamethrin, 1927
esterases, 472
fitness costs, 1927
German cockroach, 894
Heliothis virescens, 1927
horizontal transfer, 894
incompletely dominant, 472
Pieris rapae, 461
Plutella xylostella, 461
resistance, 1927
spinosad, 461
Spodoptera litura, 472
Trichoplusia ni, 461
induced defense,
host plant resistance, 1466
induced resistance, 1466
induced resistance,
host plant resistance, 1466
induced defense, 1466
infestation,
Benin, 1967
cassava chips, 1967
Prostephanus truncatus, 1967
yam chips, 1967
insect,
ethyl formate, 623
fumigation, 623
methyl isothiocyanate, 623
residue, 623
insect behavior,
Diabrotica virgifera virgifera, 341
fatty acid, 341
feeding stimulants, 341
Zea mays, 341
insect control,
atmospheric pressure plasma, 302
dielectric barrier discharge, 302
oxygenated helium plasma, 302
quarantine, 302
insect growth regulator,
Bemisia tabaci, 927
haplodiploid, 927
insecticide resistance, 927
pyriproxyfen, 927
insect infestation,
packaging, 1006
pasta, 1006
Sitophilus oryzae, 1006
Sitophilus zeamais, 1006
insect injury,
Bacillus thuringiensis corn, 555
corn hybrids, 555
western bean cutworm, 555
insect movement,
integrated pest management, 850
Stelidota geminata, 850
insect pathology, 660, 1088
alfalfa leafcutting bee, 660, 1088
bees, 660, 1088
chalkbrood, 660, 1088
disease control, 660
fungicide toxicity, 1088
insect pest,
Alphitobius diaperinus, 61
movement, 61
poultry litter, 61
Tenebrionidae, 61
insect survey,
community ecology, 998
feed mills, 998
stored-product insects, 998
trapping, 998
insect vector,
disease management, 1
epidemiology, 1
modeling, 1
insectary crops,
aphid predators, 1526
conservation biocontrol, 1526
organic agriculture, 1526
Syrphidae, 1526
insecticidal ear tags,
permethrin, 1697
piperonyl butoxide, 1697
release, 1697
resistance, 1697
insecticide, 168, 360, 391, 438, 1204, 1439, 1600
Anthonomus grandis grandis, 1600
Aphis glycines, 391
cabbage, 438
Culex, 1204
Gossypium hirsutum, 1600
Harmonia axyridis, 391
harvest, 1600
Heliothis virescens, 168
instar, 1204
LC_{50}, 1439
mosquito, 1204
Ochlerotatus, 1204
onion thrips, 438
organic, 391
plant resistance, 438
proactive, 1600
resistance management, 168
sex-related response, 1439
silkworm, 360
soybean, 391
stink bug, 1439
Thrips tabaci, 438
toxicity, 360
transgenic cotton, 168
insecticide resistance, 190, 480, 499, 927, 1685, 1897, 1919
acrinathrin, 499
Alphitobius diaperinus, 480
apple pests, 190
Bemisia tabaci, 927
cropping-structure, 1685
Cydia pomonella, 190
esterase alleles, 1897
fitness, 499
gene flow, 190
haplodiploid, 927
insect growth regulator, 927
integrated pest management, 1919
knockdown, 480
lesser mealworm, 480
microsatellites, 190
mosquitoes, 1897
Myzus persicae, 1919
peach, 1919
potato, 1919
pyriproxyfen, 927
recovery, 480

spinosad, 499
Thrips, 1685
Thysanoptera, 499
insecticide resistance genes,
Culex pipiens, 484
esterase, 484
mosquito, 484
insecticide resistance management, 492, 1406
Bt corn, 492
cornstalk borers, 492
Cry1Ac, 1406
Diatraea saccharalis, 492
Heliothis virescens, 1406
resistance monitoring, 492
single-pair family, 1406
YHD2, 1406
insecticide toxicity, 365, 375
Agriotes obscurus, 375
Elateridae, 365
fipronil, 375
morbidity, 365
neonicotinoids, 365
survival time, 375
wireworm, 375
wireworms, 365
instar,
Culex, 1204
insecticide, 1204
mosquito, 1204
Ochlerotatus, 1204
integrated pest management, 56, 850, 1302, 1594, 1864, 1919, 1950
Bacillus thuringiensis, 1950
bloom, 56
Bollgard II, 1594
bollworm, 1594, 1950
Conotrachelus nenuphar, 1302
cotton, 1950
density thresholds, 1864
Dicyphinae, 1864
dispersal, 56
Helicoverpa, 1594
insect movement, 850
insecticide resistance, 1919
Myzus persicae, 1919
omnivory, 1864
peach, 1919
pheromone, 1302
plant tolerance, 1864
plum curculio, 1302
potato, 1919
refuge, 56
Stelidota geminata, 850
tobacco budworm, 1950
weevil, 1302
Widestrike, 1594
internal transcribed spacer,
biosecurity, 1167
nuclear ribosomal RNA gene, 1167
restriction fragment length polymorphism, 1167
intrinsic rate of increase,
biotype, 955
Russian wheat aphid, 955
intrinsic rate of population increase,
mean relative growth rate, 1658
Plutella xylostella, 1658
resistance, 1658
spinosad, 1658
invasive species, 523, 720, 750, 829, 838, 1068, 1152, 1276, 1443
ant baits, 1068
ant management, 1068
bark beetle, 829
Cacopsylla chinensis, 1152
Canada thistle, 720
Cirsium arvense, 720
cytochrome oxidase I, 1152
Diaprepes abbreviatus L., 1443
distribution, 829
DNA barcoding, 523
DNA extraction, 523
eradication, 838
exotic species, 1276
floral volatiles, 720
genetic diversity, 1443
host range, 829
kairomone, 720
Lauraceae, 1276
laurel wilt, 1276
little fire ant, 1068
mitochondrial markers, 1443
Olea europaea, 750
olive fruit fly, 750
performance, 750
Pinus, 829
quarantine, 838
quarantine pest, 523, 1443
16S rDNA, 1152
restriction fragment length polymorphism, 1152
species identification, 523
Tephritidae, 750
unicolonial, 1068
IPM,
binomial data, 1719
monitoring, 1719
sequential sampling, 1719
Ipomoea batatas,
conservation tillage, 1871
no-till, 1871
pitfall traps, 1871
Ips,
Dendroctonus, 1253
multiple regression models, 1253
Pinus ponderosa, 1253
stand characteristics, 1253
Irbisia,
black grass bugs, 309
Capsus, 309
grazing management, 309
Labops, 309
Isoptera,
Coptotermes formosanus, 592
mark–recapture protocol, 592
population estimate, 592
probability density function, 592

K

kairomone, 720, 1266
Canada thistle, 720
Cirsium arvense, 720
floral volatiles, 720
invasive species, 720
mountain pine beetle, 1266
pheromones, 1266
semiochemicals, 1266
kaolin,
apples, 145
beneficials, 145
control, 145
pests, 145
knockdown,
Alphitobius diaperinus, 480
insecticide resistance, 480
lesser mealworm, 480
recovery, 480

L

Labops,
black grass bugs, 309
Capsus, 309
grazing management, 309
Irbisia, 309
landscape,
emergence, 1575
Leptinotarsa decemlineata, 1575
sampling, 1575
larval infestation,
Cydia pomonella, 81
fruit phenology, 81
monitoring, 81
within-canopy distribution, 81
Lauraceae,
exotic species, 1276
invasive species, 1276
laurel wilt, 1276
laurel wilt,
exotic species, 1276
invasive species, 1276
Lauraceae, 1276
LC_{50},
insecticide, 1439
sex-related response, 1439
stink bug, 1439

leaf toughness,
defoliation, 944
elm leaf beetle, 944
trichomes, 944
Ulmus spp., 944
leafhopper vectors,
Cicadellinae, 1114
citrus variegated chlorosis, 1114
coffee leaf scorch, 1114
leafhoppers,
deficit irrigation, 1287
grape, 1287
vineyard, 1287
water, 1287
Lepidoptera, 546, 769
attractant, 769
Bollgard, 546
Bollgard II, 546
Cry1Ac, 546
Cry2Ab2, 546
(9*E*,11*Z*)-hexadecadien-1-yl acetate, 769
pheromone antagonist, 769
pheromone race, 769
Lepinotus reticulatus,
heat shock proteins, 1974
heat stress, 1974
heat treatment, 1974
Liposcelis entomophila, 1974
Leptinotarsa decemlineata, 74, 859, 1575
aphicide, 74
eggplant, 859
emergence, 1575
landscape, 1575
potato, 859
sampling, 1575
solanaceous, 859
Solanum tuberosum, 74
tomato, 859
lesser mealworm,
Alphitobius diaperinus, 480
insecticide resistance, 480
knockdown, 480
recovery, 480
lettuce,
broccoli, 1786
Frankliniella occidentalis, 1786
phosphine, 1786
quarantine control, 1786
life history,
developmental rates, 605
marking, 605
psocids, 605
stored products, 605
life table, 9, 325, 692, 982
Anagasta kuehniella, 982
Apanteles sp., 692
biological control, 982
Brachycyrtus sp., 692
cereals, 9
development, 325
electrical penetration graph, 9
fecundity, 9, 325
Habrobracon hebetor, 982
Macrocentrus sp., 692
Myzus persicae, 9
oviposition choice tests, 692
Plodia interpunctella, 982
Scymnus subvillosus, 325
life table parameters,
Cardinium, 1711
development, 1711
Liposcelis bostrychophila, 1711
reproduction, 1711
Ligustrum sinense,
Argopistes tsekooni, 1146
flea beetle, 1146
host specificity, 1146
weed biological control, 1146
Linepithema humile, 699, 1397
Argentine ant, 699
fipronil, 1397
liquid baits, 699
necrophoresis, 1397
Pseudococcidae, 699
spot treatment, 1397
vineyards, 699
Liposcelis bostrychophila,
Cardinium, 1711
development, 1711
life table parameters, 1711
reproduction, 1711
Liposcelis entomophila,
heat shock proteins, 1974
heat stress, 1974
heat treatment, 1974
Lepinotus reticulatus, 1974
liquid baits,
Argentine ant, 699
Linepithema humile, 699
Pseudococcidae, 699
vineyards, 699
Liriomyza sativae,
biological agent, 1510
Liriomyza trifolii, 1510
multiplex polymerase chain reaction, 1510
Neochrysocharis formosa, 1510
Liriomyza trifolii,
biological agent, 1510
Liriomyza sativae, 1510
multiplex polymerase chain reaction, 1510
Neochrysocharis formosa, 1510
little fire ant,
ant baits, 1068
ant management, 1068
invasive species, 1068
unicolonial, 1068
livestock pests,
biting flies, 1034
cloth targets, 1034
fly control, 1034
low toxic,
Blattella germanica, 1384
fatty acid salts, 1384
Periplaneta americana, 1384
soap, 1384
lure,
attractant, 126
Ceratitis capitata, 126
fruit fly, 126
trap, 126
Lycoriella ingenua,
fumigant, 139
plant essential oils, 139
lysozyme,
fecundity, 777
Galleria mellonella, 777
sodium tetraborate, 777
survival, 777

M

Macrocentrus sp.,
Apanteles sp., 692
Brachycyrtus sp., 692
life table, 692
oviposition choice tests, 692
maize, 50, 90
Busseola fusca, 90
Cotesia flavipes, 50
nitrogen and potassium, 90
stemborer, 50, 90
trap plants, 50
wild hosts, 50
yield loss, 90
maize weevil,
common smut, 1455
corn earworm, 1455
husk extension, 1455
stink bugs, 1455
male annihilation,
Bactrocera cucurbitae, 759
Bactrocera dorsalis, 759
fruit fly suppression, 759
SPLAT, 759
Mamestra brassicae,
Brassica campestris subsp. *napus* variety *pekinensis*, 1134
Bt transgenic Chinese cabbage, 1134
enzyme-linked immunosorbent assay, 1134
Microplitis mediator, 1134
management,
seasonal phenology, 1847
Tomarus subtropicus, 1847
turfgrass, 1847
white grubs, 1847
mango pulp,
damage, 114
germination rate, 114

mango seeds, 114
mango weevil, 114
mango seeds,
damage, 114
germination rate, 114
mango pulp, 114
mango weevil, 114
mango weevil,
damage, 114
germination rate, 114
mango pulp, 114
mango seeds, 114
marginal vegetation,
Coccinellidae, 1792
Hippodamia variegata, 1792
supplemental food, 1792
marker-assisted selection,
corn earworm, 1471
Glycine max, 1471
quantitative trait loci, 1471
soybean looper, 1471
marking,
developmental rates, 605
life history, 605
psocids, 605
stored products, 605
mark–recapture protocol,
Coptotermes formosanus, 592
Isoptera, 592
mark–recapture protocol, 592
population estimate, 592
probability density function, 592
mass trapping,
Pheromone, 1295
population suppression, 1295
Prays nephelomima, 1295
material effect,
Conotrachelus nenuphar, 1838
monitoring, 1838
traps, 1838
weather effect, 1838
maternal and progeny fitness,
adult, 1760
cold storage, 1760
demographics, 1760
Conatocerus ashmeadi, 1760
mating behavior,
Cylas formicarius elegantulus, 1198
gamma radiation, 1198
sterile insect technique (SIT), 1198
sweetpotato weevil, 1198
mating disruption, 444, 1568, 1633, 1856
almonds, 1633
Amyelois transitella, 1633
(*E*,*Z*)-7,9-dodecadienyl acetate, 444
false-trail following, 444
granule formulation, 1856
grapevine, 444
mating rate, 1568
oriental beetle, 1856
pistachios, 1633
sex pheromone, 1568
soywax, 1856
turfgrass, 1856
wireworm, 1568
mating rate,
mating disruption, 1568
sex pheromone, 1568
wireworm, 1568
mean relative growth rate,
intrinsic rate of population increase, 1658
Plutella xylostella, 1658
resistance, 1658
spinosad, 1658
Medicago,
alfalfa leafcutting bee, 674
Apoidea, 674
Megachilidae, 674
pollen ball, 674
Megachile rotundata,
methodology, 784
Osmia lignaria, 784
pesticides, 784
sublethal effects, 784
Megachilidae,
alfalfa leafcutting bee, 674
Apoidea, 674
Medicago, 674
pollen ball, 674
Menacanthus stramineus,
humidity, 637
Ornithonyssus sylviarum, 637
survival, 637
temperature, 637
metabolism,
Chilo suppressalis, 1415
fipronil, 1415
Sesamia inferens, 1415
susceptibility, 1415
Metarhizium,
entomopathogenic fungus, 885
epizootic, 885
termite, 885
methodology,
Megachile rotundata, 784
Osmia lignaria, 784
pesticides, 784
sublethal effects, 784
methyl eugenol,
attract-and-kill system, 352
Bactrocera dorsalis, 352
neonicotinoid insecticides, 352
oriental fruit fly, 352
methyl isothiocyanate,
ethyl formate, 623
fumigation, 623
insect, 623
residue, 623
3-methylcyclohex-2-en-1-one,
antiaggregation pheromones, 1826
Dendroctonus pseudotsugae, 1826
Mexican rice borer,
movement, 237
rice, 237
sugarcane, 237
microaggregation,
Bacillus thuringiensis, 36
benzylideneacetone, 36
immunosuppression, 36
Spodoptera exigua, 36
microbial contaminants,
nucleopolyhedrovirus, 42
Spodoptera exigua, 42
storage, 42
microbial control,
biological control, 251
entomopathogenic nematodes, 251
strawberry, 251
microclimate,
augmentation biological control, 421
predation, 421
Rhyacionia frustrana, 421
Trichogramma exiguum, 421
Microplitis mediator,
Brassica campestris subsp. *napus* variety *pekinensis*, 1134
Bt transgenic Chinese cabbage, 1134
enzyme-linked immunosorbent assay, 1134
Mamestra brassicae, 1134
microsatellite molecular markers,
genotyping, 1062
plowing, 1062
Solenopsis invicta, 1062
microsatellites, 190, 1349
apple pests, 190
Cydia pomonella, 190
gene flow, 190
insecticide resistance, 190
population genetics, 1349
R. hageni, 1349
Reticulitermes flavipes, 1349
Reticulitermes virginicus, 1349
microwaves,
barley, 1011
disinfestation, 1011
mortality, 1011
stored-grain insects, 1011
migration,
fall armyworm, 742
haplotype, 742
Spodoptera frugiperda, 742

mites,
calcofluor, 1028
chitinase, 1028
diflubenzuron, 1028
peritrophic membrane, 1028
miticides,
Apis mellifera, 1081
drones, 1081
honey bee, 1081
spermatozoa, 1081
mitochondrial markers,
Diaprepes abbreviatus L., 1443
genetic diversity, 1443
invasive species, 1443
quarantine pest, 1443
model evaluation,
Anarsia lineatella, 1557
bionomics, 1557
developmental rate, 1557
temperature, 1557
modeling, 1, 989
disease management, 1
epidemiology, 1
heat treatments, 989
insect vector, 1
survival, 989
Tribolium confusum, 989
molecular diagnostic test,
Daktulosphaira vitifoliae, 1934
DNA, 1934
grapevine, 1934
pest detection, 1934
soil, 1934
monitoring, 81, 399, 404, 1719, 1821, 1838
attractant synergist, 1821
bark beetles, 1821
binomial data, 1719
boll weevil, 399
Conotrachelus nenuphar, 1838
corn pest, 404
Cydia pomonella, 81
dichlorvos, 399
endo-brevicomin, 1821
fruit phenology, 81
grandlure, 399
IPM, 1719
larval infestation, 81
material effect, 1838
pest management, 404
pheromone, 1821
sequential sampling, 1719
Striacosta albicosta, 404
trapping, 399
traps, 1838
weather effect, 1838
within-canopy distribution, 81
Monochamus clamator,
Buprestidae, 409
Cerambycidae, 409
semiochemicals, 409
Siricidae, 409
morbidity,
Elateridae, 365
insecticide toxicity, 365
neonicotinoids, 365
wireworms, 365
morphological traits,
eco-races, 1176
genetic variability, 1176
polymorphism, 1176
tasar silk worm, 1176
mortality,
barley, 1011
disinfestation, 1011
microwaves, 1011
stored-grain insects, 1011
mosquito, 484, 1204, 1897
Culex pipiens, 484
Culex, 1204
esterase, 484
esterase alleles, 1897
insecticide, 1204
insecticide resistance, 1897
insecticide resistance genes, 484
instar, 1204
Ochlerotatus, 1204
mountain pine beetle,
kairomones, 1266
pheromones, 1266
semiochemicals, 1266
movement, 61, 237
Alphitobius diaperinus, 61
insect pest, 61
Mexican rice borer, 237
poultry litter, 61
rice, 237
sugarcane, 237
Tenebrionidae, 61
multiple regression models,
Dendroctonus, 1253
Ips, 1253
Pinus ponderosa, 1253
stand characteristics, 1253
multiple resistance,
acaricide resistance, 1704
cross-resistance, 1704
hitchhiking effect, 1704
Tetranychus urticae, 1704
multiple species approach,
Heterorhabditis bacteriophora, 1533
Otiorhynchus ligustici, 1533
Steinernema carpocapsae, 1533
Steinernema feltiae, 1533
multiple-funnel trap,
Cerambycidae, 107
Curculionidae, 107
detection, 107
Scolytidae, 107
multiplex PCR,
armored scale insects, 1944
cytochrome oxidase I and *II* genes, 1944
distribution and species composition, 1944
DNA diagnostic, 1944
multiplex polymerase chain reaction,
biological agent, 1510
Liriomyza sativae, 1510
Liriomyza trifolii, 1510
Neochrysocharis formosa, 1510
Myzus persicae, 9, 728, 1432, 1919
cereals, 9
colonization behavior, 728
electrical penetration graph, 9
electronic penetration graph, 728
fecundity, 9
fertility life table, 1432
fitness cost, 1432
insecticide resistance, 1919
integrated pest management, 1919
life table, 9
peach, 1919
potato, 1919
probing behavior, 728
resistance to insecticide, 1432
Rhopalosiphum padi, 728

N

natural fumigant,
botanical insecticide, 1960
cinnamaldehyde, 1960
Sitophilus oryzae, 1960
structure–activity relationship, 1960
necrophoresis,
fipronil, 1397
Linepithema humile, 1397
spot treatment, 1397
Neochrysocharis formosa,
biological agent, 1510
Liriomyza sativae, 1510
Liriomyza trifolii, 1510
multiplex polymerase chain reaction, 1510
neonicotinoid, 159, 365, 451, 1881
California red scale, 451
canola, 159
citricola scale, 451
citrus red mite, 451
cottony cushion scale, 451
Elateridae, 365
flea beetles, 159
insecticide toxicity, 365
morbidity, 365
seed treatment, 159
soil penetration, 1881
species composition shift, 159

termite mortality, 1881
treatment depth, 1881
wireworms, 365
neonicotinoid insecticides,
attract-and-kill system, 352
Bactrocera dorsalis, 352
methyl eugenol, 352
oriental fruit fly, 352
Neozygites fresenii,
Aphis gossypii, 23
biological control, 23
epizootics, 23
Nilaparvata lugens, 384, 515, 575
assimilate allocation, 384
genetics, 575
imidacloprid, 515
physiological response, 384
resistance, 515, 575
rice, 384
rice mutants, 575
risk assessment, 515
susceptibilities, 515
nitrogen and potassium,
Busseola fusca, 90
maize, 90
stemborer, 90
yield loss, 90
no-till,
conservation tillage, 1871
Ipomoea batatas, 1871
pitfall traps, 1871
Nosema spp.,
Acarapis woodi, 668
Apis mellifera, 668
package honey bees, 668
Varroa destructor, 668
noviflumuron,
Coptotermes formosanus, 880
feeding preference, 880
gamma-irradiated wood, 880
hexaflumuron, 880
nuclear ribosomal RNA gene,
biosecurity, 1167
internal transcribed spacer, 1167
restriction fragment length polymorphism, 1167
nucleopolyhedrovirus,
microbial contaminants, 42
Spodoptera exigua, 42
storage, 42
nursery,
Popillia japonica, 1158
quarantine, 1158
scarab, 1158
white grubs, 1158
nutrition,
Aphis glycines, 1228
artificial diet, 1228
soybean aphid, 1228
nutritional ecology,
Diabrotica virgifera virgifera, 1547
plant–insect interactions, 1547
nutritive value,
Bombus terrestris, 1737
comparative bioassay, 1737
pollen consumption, 1737
pollen diet, 1737

O

Oberon,
Bemisia argentifolii, 174
imidacloprid-resistant whiteflies, 174
Q Biotype, 174
whitefly immature stages, 174
Ochlerotatus,
Culex, 1204
insecticide, 1204
instar, 1204
mosquito, 1204
Oebalus pugnax, 216, 1233
attractiveness, 1233
Oryza sativa, 216, 1233
rice panicle stage, 1233
rice stink bug, 216, 1233
sweep net sampling, 216
visual sampling, 216
Olea europaea,
invasive species, 750
olive fruit fly, 750
performance, 750
Tephritidae, 750
olfactometry,
electrical penetration graph, 584
feeding behavior, 584
Solanum chomatophilum, 584
Solanum stoloniferum, 584
olive fruit fly,
invasive species, 750
Olea europaea, 750
performance, 750
Tephritidae, 750
omnivory,
density thresholds, 1864
Dicyphinae, 1864
integrated pest management, 1864
plant tolerance, 1864
onion thrips, 438, 1331
Allium cepa, 1331
cabbage, 438
insecticide, 438
plant resistance, 438
Thrips tabaci, 438, 1331
within-plant distribution, 1331
onions,
cultural management, 1317
habitat management, 1317
straw mulch, 1317
Thrips tabaci, 1317
orange trees,
glassy-winged sharpshooter, 226
pest management, 226
sex ratio, 226
yellow sticky trap, 226
orchard pollination,
behavioral effects, 647
side effects, 647
sublethal dosages, 647
organic,
Aphis glycines, 391
Harmonia axyridis, 391
insecticides, 391
soybean, 391
organic agriculture,
aphid predators, 1526
conservation biocontrol, 1526
insectary crops, 1526
Syrphidae, 1526
organophosphates, 1667, 1676
carbamates, 1667
Chrysoperla carnea, 1676
pyrethroids, 1667, 1676
resistance, 1667, 1676
Spodoptera litura, 1667
stability of resistance, 1676
oriental beetle,
granule formulation, 1856
mating disruption, 1856
soywax, 1856
turfgrass, 1856
oriental fruit fly,
attract-and-kill system, 352
Bactrocera dorsalis, 352
methyl eugenol, 352
neonicotinoid insecticides, 352
oriental fruit moth,
CATTS, 295
codling moth, 295
controlled atmospheres, 295
heat, 295
Orius,
beat bucket, 1501
beneficial arthropods, 1501
cotton, 1501
spiders, 1501
Orius laevigatus,
artificial diet, 1127
fecundity, 1127
quality assurance, 1127
Ornithonyssus sylviarum,
humidity, 637
Menacanthus stramineus, 637
survival, 637
temperature, 637
Orosius orientalis,
Circulifer spp., 1337
Exitianus capicola, 1337
physical control, 1337
UV-absorbing plastic, 1337

Oryza sativa, 216, 1233
attractiveness, 1233
Oebalus pugnax, 216, 1233
rice panicle stage, 1233
rice stink bug, 216, 1233
sweep net sampling, 216
visual sampling, 216
Osmia lignaria,
Megachile rotundata, 784
methodology, 784
pesticides, 784
sublethal effects, 784
Ostrinia nubilalis,
Bt corn, 1614
European corn borer, 1614
remote sensing, 1614
resistance, 1614
Otiorhynchus ligustici,
Heterorhabditis bacteriophora, 1533
multiple species approach, 1533
Steinernema carpocapsae, 1533
Steinernema feltiae, 1533
oviposition,
behavior, 118
particle film technology, 118
Rhagoletis mendax, 118
Surround WP, 118
oviposition choice tests,
Apanteles sp., 692
Brachycyrtus sp., 692
life table, 692
Macrocentrus sp., 692
oxalic acid,
Apis mellifera, 1057
desert conditions, 1057
Sucrocide, 1057
Varroa destructor, 1057
oxygenated helium plasma,
atmospheric pressure plasma, 302
dielectric barrier discharge, 302
insect control, 302
quarantine, 302

P

package honey bees,
Acarapis woodi, 668
Apis mellifera, 668
Nosema spp., 668
Varroa destructor, 668
packaging,
insect infestation, 1006
pasta, 1006
Sitophilus oryzae, 1006
Sitophilus zeamais, 1006
Paenibacillus larvae,
American foulbrood disease, 1095
Apis mellifera, 1095
shaking technique, 1095
shook-swarm, 1095
pan evaporation,
climatic conditions, 334
Dermolepida albohirtum, 334
sugarcane, 334
time series, 334
parasitic Hymenoptera,
biological control, 1140
sex determination, 1140
sex ratio, 1140
parasitoid,
Aphis glycines, 278
classical biological control, 278
ethylene glycol, 631
Plodia interpunctella, 631
predatory flies, 278
Sitophilus spp., 631
trapping, 631
paratransgenesis,
Alcaligenes xylosoxidans, 1519
arthropod vector, 1519
Pierce's disease, 1519
symbiotic control, 1519
particle film technology,
behavior, 118
oviposition, 118
Rhagoletis mendax, 118
Surround WP, 118
pasta,
insect infestation, 1006
packaging, 1006
Sitophilus oryzae, 1006
Sitophilus zeamais, 1006
pea,
adult colonization, 1606
pea leaf weevil, 1606
reduced tillage, 1606
pea leaf weevil,
adult colonization, 1606
pea, 1606
reduced tillage, 1606
peach,
insecticide resistance, 1919
integrated pest management, 1919
Myzus persicae, 1919
potato, 1919
Pectinophora gossypiella,
Bt cotton, 504
pink bollworm, 504
refuge, 504
resistance, 504
performance,
invasive species, 750
Olea europaea, 750
olive fruit fly, 750
Tephritidae, 750
Periplaneta americana,
Blattella germanica, 1384
fatty acid salts, 1384
low toxic, 1384
soap, 1384
peritrophic membrane,
calcofluor, 1028
chitinase, 1028
diflubenzuron, 1028
mites, 1028
permethrin,
insecticidal ear tags, 1697
piperonyl butoxide, 1697
release, 1697
resistance, 1697
Persea americana,
avocado pests, 1310
Braconidae, 1310
Ichneumonidae, 1310
quarantine, 1310
pest detection,
Daktulosphaira vitifoliae, 1934
DNA, 1934
grapevine, 1934
molecular diagnostic test, 1934
soil, 1934
pest management, 226, 404, 810, 1211
canola, 810
Cerotoma trifurcata, 1211
corn pest, 404
cultural control, 810
flea beetles, 810
glassy-winged sharpshooter, 226
Glycine max, 1211
monitoring, 404
orange trees, 226
Phyllotreta spp., 810
seed treatment, 1211
sex ratio, 226
Striacosta albicosta, 404
yellow sticky trap, 226
pesticide residue,
baculovirus, 1540
biopesticide, 1540
PoGV, 1540
Solanum tuberosum, 1540
pesticides,
Megachile rotundata, 784
methodology, 784
Osmia lignaria, 784
sublethal effects, 784
pests,
apples, 145
beneficials, 145
control, 145
kaolin, 145
phenology,
biofix, 1651
codling moth, 1651
pheromone, 1266, 1295, 1302, 1325, 1821
attractant synergist, 1821
bark beetles, 1821
Ceratitis capitata, 1325

Conotrachelus nenuphar, 1302
dispenser, 1325
endo-brevicomin, 1821
fruit fly, 1325
integrated pest management, 1302
kairomones, 1266
mass trapping, 1295
monitoring, 1821
mountain pine beetle, 1266
plum curculio, 1302
population suppression, 1295
Prays nephelomima, 1295
semiochemicals, 1266
trimedlure, 1325
weevil, 1302
pheromone antagonist,
attractant, 769
(9*E*,11Z)-hexadecadien-1-yl acetate, 769
Lepidoptera, 769
pheromone race, 769
pheromone race,
attractant, 769
(9*E*,11Z)-hexadecadien-1-yl acetate, 769
Lepidoptera, 769
pheromone antagonist, 769
Phoracantha,
biological control, 1771
collection strategy, 1771
egg parasitoid, 1771
eucalyptus longhorned borer, 1771
phosphine,
broccoli, 1786
Frankliniella occidentalis, 1786
lettuce, 1786
quarantine control, 1786
Phyllophaga,
Cynodon, 1624
cytochrome c oxidase I, 1624
scarab flight periods, 1624
white grub, 1624
Phyllotreta spp., 810, 1811
canola, 810, 1811
cultural control, 810
decreased pesticide load, 1811
flea beetle injury, 1811
flea beetles, 810
pest management, 810
seed treatments, 1811
physical control,
Circulifer spp., 1337
Exitianus capicola, 1337
Orosius orientalis, 1337
UV-absorbing plastic, 1337
physiological response,
assimilate allocation, 384
Nilaparvata lugens, 384
rice, 384
physiological time model,
corn rootworm, 1584
phytosanitary treatment,
Amorbia emigratella, 710
postharvest, 710
quarantine pest, 710
Pierce's disease, 1041, 1122, 1519
Alcaligenes xylosoxidans, 1519
arthropod vector, 1519
epidemiology, 1041
Hemiptera, 1122
Homalodisca vitripennis, 1041
paratransgenesis, 1519
Pierce's disease, 1041, 1122, 1519
quantification, 1122
real-time polymerase chain reaction, 1122
symbiotic control, 1519
transmission, 1122
Xylella fastidiosa, 1041
Pierce's disease (PD) of grapevines,
almond leaf scorch, 1105
grass sharpshooter, 1105
Pierce's disease (PD) of grapevines, 1105
population dynamics, 1105
transmission efficiency, 1105
Pieris rapae,
indoxacarb, 461
Plutella xylostella, 461
spinosad, 461
Trichoplusia ni, 461
pink bollworm,
Bt cotton, 504
Pectinophora gossypiella, 504
refuge, 504
resistance, 504
Pinus,
bark beetle, 829
distribution, 829
host range, 829
invasive species, 829
Pinus ponderosa,
Dendroctonus, 1253
Ips, 1253
multiple regression models, 1253
stand characteristics, 1253
piperonyl butoxide,
insecticidal ear tags, 1697
permethrin, 1697
release, 1697
resistance, 1697
pistachios,
almonds, 1633
Amyelois transitella, 1633
mating disruption, 1633
pitfall traps,
conservation tillage, 1871
Ipomoea batatas, 1871
no-till, 1871
plant essential oils,
fumigant, 139
Lycoriella ingenua, 139
plant phenology,
Aphis glycines, 1488
coefficient of variation, 1488
sampling, 1488
plant resistance, 438, 533, 569
biotype, 569
Blissus occiduus, 533
cabbage, 438
chinch bug, 533
Diuraphis noxia, 569
insecticide, 438
onion thrips, 438
Thrips tabaci, 438
Triticum aestivum, 569
turfgrass, 533
plant tolerance,
density thresholds, 1864
Dicyphinae, 1864
integrated pest management, 1864
omnivory, 1864
planting date,
Chaetocnema confinis, 1238
harvest date, 1238
soil temperature, 1238
sweetpotato, 1238
plant–insect interactions,
Diabrotica virgifera virgifera, 1547
nutritional ecology, 1547
plant–insect interactions, 1547
Plodia interpunctella, 631, 982, 1051
Anagasta kuehniella, 982
biological control, 982
commodity treatment, 1051
ethylene glycol, 631
Habrobracon hebetor, 982
life tables, 982
parasitoid, 631
quarantine, 1051
radiation, 1051
Sitophilus spp., 631
Sitotroga cerealella, 1051
trapping, 631
plowing,
genotyping, 1062
microsatellite molecular markers, 1062
Solenopsis invicta, 1062
plum curculio,
Conotrachelus nenuphar, 1302
integrated pest management, 1302
pheromone, 1302
weevil, 1302

Plutella xylostella, 461, 1658, 1888
abamectin, 1888
functional expression, 1888
GABA receptor, 1888
indoxacarb, 461
intrinsic rate of population increase, 1658
mean relative growth rate, 1658
Pieris rapae, 461
resistance, 1658
spinosad, 461, 1658
Trichoplusia ni, 461
Xenopus oocyte, 1888
PoGV,
baculovirus, 1540
biopesticide, 1540
pesticide residue, 1540
Solanum tuberosum, 1540
pollen ball,
alfalfa leafcutting bee, 674
Apoidea, 674
Medicago, 674
Megachilidae, 674
pollen consumption,
Bombus terrestris, 1737
comparative bioassay, 1737
nutritive value, 1737
pollen diet, 1737
pollen diet,
Bombus terrestris, 1737
comparative bioassay, 1737
nutritive value, 1737
pollen consumption, 1737
polymorphism,
eco-races, 1176
genetic variability, 1176
morphological traits, 1176
tasar silk worm, 1176
Popillia japonica,
nursery, 1158
quarantine, 1158
scarab, 1158
white grubs, 1158
population dynamics,
almond leaf scorch, 1105
grass sharpshooter, 1105
Pierce's disease (PD) of grapevines, 1105
transmission efficiency, 1105
population estimate,
Coptotermes formosanus, 592
Isoptera, 592
mark–recapture protocol, 592
probability density function, 592
population genetics, 272, 1349, 1729
Anthonomus grandis grandis, 1729
Apis, 272
bee breeding, 272
boll weevil, 1729
dispersal, 1729
eradication, 1729
feral honey bees, 272
microsatellites, 1349
R. hageni, 1349
Reticulitermes flavipes, 1349
Reticulitermes virginicus, 1349
population suppression,
mass trapping, 1295
Pheromone, 1295
Prays nephelomima, 1295
Populus maximowiczii,
Cryptorhynchus lapathi, 199
host plant resistance, 199
hybrid poplars, 199
postharvest,
Amorbia emigratella, 710
phytosanitary treatment, 710
quarantine pest, 710
postharvest pest control,
black widow spider, 1515
controlled atmosphere, 1515
table grape, 1515
ultralow oxygen, 1515
potato, 859, 1919
eggplant, 859
insecticide resistance, 1919
integrated pest management, 1919
Leptinotarsa decemlineata, 859
Myzus persicae, 1919
peach, 1919
solanaceous, 859
tomato, 859
potato leafhopper,
Bt cotton, 959
economic threshold, 959
F_2 screen, 959
host plant resistance, 959
resistance detection, 959
sap-feeding insect, 959
tolerance, 959
poultry litter,
Alphitobius diaperinus, 61
insect pest, 61
movement, 61
Tenebrionidae, 61
Prays nephelomima,
mass trapping, 1295
Pheromone, 1295
population suppression, 1295
predation,
augmentation biological control, 421
microclimate, 421
Rhyacionia frustrana, 421
Trichogramma exiguum, 421
predatory flies,
Aphis glycines, 278
classical biological control, 278
parasitoids, 278
prism trap,
emerald ash borer, 1831
purple, 1831
trap height, 1831
visual cues, 1831
proactive,
Anthonomus grandis grandis, 1600
Gossypium hirsutum, 1600
harvest, 1600
insecticide, 1600
probability density function,
Coptotermes formosanus, 592
Isoptera, 592
mark–recapture protocol, 592
population estimate, 592
probing,
electrical penetration graph (EPG), 1183
Homalodisca coagulata, 1183
host plant acceptance, 1183
probing behavior,
colonization behavior, 728
electronic penetration graph, 728
Myzus persicae, 728
Rhopalosiphum padi, 728
Prostephanus truncatus,
Benin, 1967
cassava chips, 1967
infestation, 1967
yam chips, 1967
Pseudococcidae,
Argentine ant, 699
Linepithema humile, 699
liquid baits, 699
vineyards, 699
pseudoscent,
Blattella germanica, 1389
Camponotus floridanus, 1389
Cimex lectularius, 1389
dog, 1389
psocids,
developmental rates, 605
life history, 605
marking, 605
stored products, 605
Psocoptera,
artificial diets, 616
booklice, 616
sex ratio, 616
stored products, 616
puparial morphology,
Bactrocera dorsalis, 736
eclosion time, 736
heat exposure, 736
timing of death, 736
purple,
emerald ash borer, 1831
prism trap, 1831
trap height, 1831
visual cues, 1831

PVC strips,
Apis mellifera, 1075
residues, 1075
rotenone, 1075
Varroa destructor, 1075
pyrethroids, 1667, 1676, 1911
carbamates, 1667
Chrysoperla carnea, 1676
cross-resistance, 1911
Cry1Ac, 1911
diamondback moth, 1911
genetics of resistance, 1911
organophosphates, 1667, 1676
resistance, 1667, 1676
Spodoptera litura, 1667
stability of resistance, 1676
pyriproxyfen,
Bemisia tabaci, 927
haplodiploid, 927
insect growth regulator, 927
insecticide resistance, 927

Q

Q Biotype,
Bemisia argentifolii, 174
imidacloprid-resistant whiteflies, 174
Oberon, 174
whitefly immature stages, 174
quality assurance,
artificial diet, 1127
fecundity, 1127
Orius laevigatus, 1127
quantification,
Hemiptera, 1122
Pierce's disease, 1122
real-time polymerase chain reaction, 1122
transmission, 1122
quantitative trait loci, 1341, 1471
antibiosis, 1341
Aphis pomi, 1341
apple, 1341
corn earworm, 1471
Glycine max, 1471
marker-assisted selection, 1471
shoot growth characteristics, 1341
soybean looper, 1471
quarantine, 302, 716, 838, 1051, 1158, 1310, 1779
Anastrepha ludens, 716
atmospheric pressure plasma, 302
avocado pests, 1310
Braconidae, 1310
Cicadellidae, 1779
clean stock program, 1779
commodity treatment, 716, 1051
Diaspididae, 1779
dielectric barrier discharge, 302
disinfestation, 716
eradication, 838
Ichneumonidae, 1310
insect control, 302
invasive species, 838
nursery, 1158
oxygenated helium plasma, 302
Persea americana, 1310
Plodia interpunctella, 1051
Popillia japonica, 1158
radiation, 716, 1051
scarab, 1158
Sitotroga cerealella, 1051
Tettigoniidae, 1779
white grubs, 1158
quarantine control,
broccoli, 1786
Frankliniella occidentalis, 1786
lettuce, 1786
phosphine, 1786
quarantine pest, 523, 710, 1443
Amorbia emigratella, 710
Diaprepes abbreviatus L., 1443
DNA barcoding, 523
DNA extraction, 523
genetic diversity, 1443
invasive species, 523, 1443
mitochondrial markers, 1443
phytosanitary treatment, 710
postharvest, 710
species identification, 523

R

R. hageni,
microsatellites, 1349
population genetics, 1349
Reticulitermes flavipes, 1349
Reticulitermes virginicus, 1349
radiation, 716, 1051
Anastrepha ludens, 716
commodity treatment, 716, 1051
disinfestation, 716
Plodia interpunctella, 1051
quarantine, 716, 1051
16S rDNA,
Cacopsylla chinensis, 1152
cytochrome oxidase I, 1152
invasive species, 1152
restriction fragment length polymorphism, 1152
Sitotroga cerealella, 1051
real-time polymerase chain reaction,
Hemiptera, 1122
Pierce's disease, 1122
quantification, 1122
transmission, 1122
recovery,
Alphitobius diaperinus, 480
insecticide resistance, 480
knockdown, 480
lesser mealworm, 480
reduced tillage,
adult colonization, 1606
pea, 1606
pea leaf weevil, 1606
refuge, 56, 504, 915
biotechnology, 915
bloom, 56
Bt cotton, 504, 915
dispersal, 56
integrated pest management, 56
Pectinophora gossypiella, 504
pink bollworm, 504
refugia, 915
resistance, 504, 915
refugia,
biotechnology, 915
Bt cotton, 915
refuge, 915
resistance, 915
release,
insecticidal ear tags, 1697
permethrin, 1697
piperonyl butoxide, 1697
resistance, 1697
remote sensing,
Bt corn, 1614
European corn borer, 1614
Ostrinia nubilalis, 1614
resistance, 1614
Reoviridae,
Delphacidae, 17
Saccharum, 17
sugarcane, 17
repellency,
Reticulitermes hesperus, 1360
soil treatments, 1360
termiticide, 1360
repellent,
hybrid, 265
imported fire ants, 265
Solenopsis invicta Buren, 265
Solenopsis richteri, 265
reproduction, 541, 1711, 1798
Cardinium, 1711
development, 541, 1711, 1798
development thresholds, 1798
life table parameters, 1711
Liposcelis bostrychophila, 1711
Russian wheat aphid, 541
survival, 1798
temperature, 541
thermal constants, 1798
residue, 623, 1075
Apis mellifera, 1075
ethyl formate, 623
fumigation, 623

insect, 623
methyl isothiocyanate, 623
PVC strips, 1075
rotenone, 1075
Varroa destructor, 1075
resistance, 504, 515, 564, 575, 915, 1614, 1658, 1667, 1676, 1697, 1903, 1927
antibiosis, 564
biotechnology, 915
Brachiaria, 564
Bt corn, 1614
Bt cotton, 504, 915
carbamates, 1667
Chrysoperla carnea, 1676
deltamethrin, 1927
Diatraea saccharalis, 1903
European corn borer, 1614
fitness costs, 1927
genetics, 575
Heliothis virescens, 1927
imidacloprid, 515
Indoxacarb, 1927
insecticidal ear tags, 1697
intrinsic rate of population increase, 1658
mean relative growth rate, 1658
Nilaparvata lugens, 515, 575
organophosphates, 1667, 1676
Ostrinia nubilalis, 1614
Pectinophora gossypiella, 504
permethrin, 1697
pink bollworm, 504
piperonyl butoxide, 1697
Plutella xylostella, 1658
pyrethroids, 1667, 1676
refuge, 504, 915
refugia, 915
release, 1697
remote sensing, 1614
rice mutants, 575
risk assessment, 515
selection, 1903
spinosad, 1658
spittlebugs, 564
Spodoptera litura, 1667
stability of resistance, 1676
sublethal effects, 564
sugarcane borer, 1903
susceptibilities, 515
tebufenozide, 1903
resistance detection, 933, 959
Bt cotton, 933, 959
economic threshold, 959
F_1 screen, 933, 959
Helicoverpa armigera, 933
host plant resistance, 959
potato leafhopper, 959
resistance management, 933
sap-feeding insect, 959
tolerance, 959
resistance management, 168, 933, 959
Bt cotton, 933, 959
economic threshold, 959
F_1 screen, 933, 959
Helicoverpa armigera, 933
Heliothis virescens, 168
host plant resistance, 959
insecticide, 168
potato leafhopper, 959
sap-feeding insect, 959
tolerance, 959
resistance detection, 933, 959
transgenic cotton, 168
resistance monitoring,
Bt corn, 492
cornstalk borers, 492
Diatraea saccharalis, 492
insecticide resistance management, 492
resistance to insecticide,
fertility life table, 1432
fitness cost, 1432
Myzus persicae, 1432
restriction fragment length polymorphism, 1152, 1167
biosecurity, 1167
Cacopsylla chinensis, 1152
cytochrome oxidase I, 1152
internal transcribed spacer, 1167
invasive species, 1152
nuclear ribosomal RNA gene, 1167
16S rDNA, 1152
Reticulitermes flavipes, 902, 1349
horizontal transfer, 902
microsatellites, 1349
population genetics, 1349
R. hageni, 1349
Reticulitermes virginicus, 1349
termiticide, 902
Reticulitermes hesperus,
repellency, 1360
soil treatments, 1360
termiticide, 1360
Reticulitermes virginicus,
microsatellites, 1349
population genetics, 1349
R. hageni, 1349
Reticulitermes flavipes, 1349
Rhagoletis mendax,
behavior, 118
oviposition, 118
particle film technology, 118
Surround WP, 118
Rhagoletis pomonella,
apple, 206
germplasm, 206
host plant resistance, 206
Rhopalosiphum padi,
colonization behavior, 728
electronic penetration graph, 728
Myzus persicae, 728
probing behavior, 728
Rhyacionia frustrana,
augmentation biological control, 421
microclimate, 421
predation, 421
Trichogramma exiguum, 421
Rhyzopertha dominica,
grains, 976
Sitophilus oryzae, 976
spinosad, 976
rice, 237, 384
assimilate allocation, 384
Mexican rice borer, 237
movement, 237
Nilaparvata lugens, 384
physiological response, 384
sugarcane, 237
rice mutants,
genetics, 575
Nilaparvata lugens, 575
resistance, 575
rice panicle stage,
attractiveness, 1233
Oebalus pugnax, 1233
Oryza sativa, 1233
rice stink bug, 1233
rice stink bug, 216, 1233
attractiveness, 1233
Oebalus pugnax, 216, 1233
Oryza sativa, 216, 1233
rice panicle stage, 1233
sweep net sampling, 216
visual sampling, 216
risk assessment,
imidacloprid, 515
Nilaparvata lugens, 515
resistance, 515
susceptibilities, 515
rotenone,
Apis mellifera, 1075
PVC strips, 1075
residues, 1075
Varroa destructor, 1075
Russian bees,
Acarapis woodi, 654
Apis mellifera, 654
Varroa destructor, 654
varroa sensitive hygiene, 654
Russian wheat aphid, 541, 955
biotype, 955
development, 541
intrinsic rate of increase, 955
reproduction, 541
temperature, 541

S

Saccharum,
Delphacidae, 17
Reoviridae, 17
sugarcane, 17
sample size,
Diaphorina citri, 1478
dispersion, 1478
sampling units, 1478
sampling, 969, 1488, 1575
Aphis glycines, 1488
Cochylis hospes, 969
coefficient of variation, 1488
economic distance, 969
EIL, 969
emergence, 1575
Helianthus annuus, 969
landscape, 1575
Leptinotarsa decemlineata, 1575
plant phenology, 1488
sampling units,
Diaphorina citri, 1478
dispersion, 1478
sample size, 1478
sap-feeding i*Helicoverpa armigera*,
Bt cotton, 959
economic threshold, 959
F_1 screen, 959
host plant resistance, 959
potato leafhopper, 959
resistance detection, 959
tolerance, 959
scarab,
nursery, 1158
Popillia japonica, 1158
quarantine, 1158
white grubs, 1158
scarab flight periods,
Cynodon, 1624
cytochrome c oxidase I, 1624
Phyllophaga, 1624
white grub, 1624
Scolytidae,
Cerambycidae, 107
Curculionidae, 107
detection, 107
multiple-funnel trap, 107
Scymnus subvillosus,
development, 325
fecundity, 325
life table, 325
seasonal phenology,
management, 1847
Tomarus subtropicus, 1847
turfgrass, 1847
white grubs, 1847
seed treatment, 159, 1211, 1811
canola, 159, 1811
Cerotoma trifurcata, 1211
decreased pesticide load, 1811
flea beetle injury, 1811
flea beetles, 159
Glycine max, 1211
neonicotinoid, 159
pest management, 1211
species composition shift, 159
Phyllotreta spp., 1811
selection,
Diatraea saccharalis, 1903
resistance, 1903
sugarcane borer, 1903
tebufenozide, 1903
Semanotus ligneus ligneus,
Atimia confusa confusa, 430
Callidiellum rufipenne, 430
Callidium frigidum, 430
trap log, 430
semiochemicals, 409, 1266
Buprestidae, 409
Cerambycidae, 409
kairomones, 1266
Monochamus clamator, 409
mountain pine beetle, 1266
pheromones, 1266
Siricidae, 409
sequential sampling,
binomial data, 1719
IPM, 1719
monitoring, 1719
Sesamia inferens,
Chilo suppressalis, 1415
fipronil, 1415
metabolism, 1415
susceptibility, 1415
sex determination,
biological control, 1140
parasitic Hymenoptera, 1140
sex ratio, 1140
sex pheromone,
mating disruption, 1568
mating rate, 1568
wireworm, 1568
sex ratio, 226, 616, 1140
artificial diets, 616
biological control, 1140
booklice, 616
glassy-winged sharpshooter, 226
orange trees, 226
parasitic Hymenoptera, 1140
pest management, 226
Psocoptera, 616
sex determination, 1140
stored products, 616
yellow sticky trap, 226
sex-related response,
insecticide, 1439
LC_{50}, 1439
stink bug, 1439
shaking technique,
American foulbrood disease, 1095
Apis mellifera, 1095
Paenibacillus larvae, 1095
shook-swarm, 1095
shelf life,
Cydia pomonella, 288
granulovirus, 288
storage, 288
temperature, 288
shook-swarm,
American foulbrood disease, 1095
Apis mellifera, 1095
Paenibacillus larvae, 1095
shaking technique, 1095
shoot growth characteristics,
antibiosis, 1341
Aphis pomi, 1341
apple, 1341
quantitative trait loci, 1341
side effects,
behavioral effects, 647
orchard pollination, 647
sublethal dosages, 647
silkworm,
insecticide, 360
toxicity, 360
single-pair family,
Cry1Ac, 1406
Heliothis virescens, 1406
insecticide resistance management, 1406
YHD2, 1406
Siricidae,
Buprestidae, 409
Cerambycidae, 409
Monochamus clamator, 409
semiochemicals, 409
Sitophilus oryzae, 976, 1006, 1960
botanical insecticide, 1960
cinnamaldehyde, 1960
grains, 976
insect infestation, 1006
natural fumigant, 1960
packaging, 1006
pasta, 1006
Rhyzopertha dominica, 976
Sitophilus zeamais, 1006
spinosad, 976
structure–activity relationship, 1960
Sitophilus zeamais,
insect infestation, 1006
packaging, 1006
pasta, 1006
Sitophilus oryzae, 1006
Sitophilus spp.,
ethylene glycol, 631
parasitoid, 631
Plodia interpunctella, 631
trapping, 631

Sitotroga cerealella,
commodity treatment, 1051
Plodia interpunctella, 1051
quarantine, 1051
radiation, 1051
soap,
Blattella germanica, 1384
fatty acid salts, 1384
low toxic, 1384
Periplaneta americana, 1384
sodium tetraborate,
fecundity, 777
Galleria mellonella, 777
lysozyme, 777
survival, 777
soil,
Daktulosphaira vitifoliae, 1934
DNA, 1934
grapevine, 1934
molecular diagnostic test, 1934
pest detection, 1934
soil penetration,
neonicotinoids, 1881
termite mortality, 1881
treatment depth, 1881
soil temperature,
Chaetocnema confinis, 1238
harvest date, 1238
planting date, 1238
sweetpotato, 1238
soil treatments,
repellency, 1360
Reticulitermes hesperus, 1360
termiticide, 1360
solanaceous,
eggplant, 859
Leptinotarsa decemlineata, 859
potato, 859
tomato, 859
Solanum chomatophilum,
electrical penetration graph, 584
feeding behavior, 584
olfactometry, 584
Solanum stoloniferum, 584
Solanum stoloniferum,
electrical penetration graph, 584
feeding behavior, 584
olfactometry, 584
Solanum chomatophilum, 584
Solanum tuberosum, 74, 1540
aphicide, 74
baculovirus, 1540
biopesticide, 1540
Leptinotarsa decemlineata, 74
pesticide residue, 1540
PoGV, 1540
Solenopsis invicta,
genotyping, 1062
microsatellite molecular markers, 1062
plowing, 1062
Solenopsis invicta Buren,
hybrid, 265
imported fire ants, 265
repellent, 265
Solenopsis richteri, 265
Solenopsis richteri,
hybrid, 265
imported fire ants, 265
repellent, 265
Solenopsis invicta Buren, 265
soybean,
Aphis glycines, 391
Harmonia axyridis, 391
insecticides, 391
organic, 391
soybean aphid,
Aphis glycines, 1228
artificial diet, 1228
nutrition, 1228
soybean looper,
corn earworm, 1471
Glycine max, 1471
marker-assisted selection, 1471
quantitative trait loci, 1471
soywax,
granule formulation, 1856
mating disruption, 1856
oriental beetle, 1856
turfgrass, 1856
species composition shift,
canola, 159
flea beetles, 159
neonicotinoid, 159
seed treatment, 159
species identification,
DNA barcoding, 523
DNA extraction, 523
invasive species, 523
quarantine pest, 523
spermatozoa,
Apis mellifera, 1081
drones, 1081
honey bee, 1081
miticides, 1081
spiders,
beat bucket, 1501
beneficial arthropods, 1501
cotton, 1501
Orius, 1501
spinosad, 461, 499, 976, 1022, 1658
acrinathrin, 499
fitness, 499
grains, 976
indoxacarb, 461
insecticide resistance, 499
intrinsic rate of population increase, 1658
mean relative growth rate, 1658
Pieris rapae, 461
Plutella xylostella, 461, 1658
resistance, 1658
Rhyzopertha dominica, 976
Sitophilus oryzae, 976
stored-grain insects, 1022
susceptibility differences, 1022
Thysanoptera, 499
Trichoplusia ni, 461
spittlebugs,
antibiosis, 564
Brachiaria, 564
resistance, 564
sublethal effects, 564
SPLAT,
Bactrocera cucurbitae, 759
Bactrocera dorsalis, 759
fruit fly suppression, 759
male annihilation, 759
Spodoptera exigua, 36, 42
Bacillus thuringiensis, 36
benzylideneacetone, 36
immunosuppression, 36
microaggregation, 36
microbial contaminants, 42
nucleopolyhedrovirus, 42
storage, 42
Spodoptera frugiperda,
fall armyworm, 742
haplotype, 742
migration, 742
Spodoptera litura, 472, 1667
autosomal, 472
carbamates, 1667
esterases, 472
incompletely dominant, 472
indoxacarb, 472
organophosphates, 1667
pyrethroids, 1667
resistance, 1667
spot treatment,
fipronil, 1397
Linepithema humile, 1397
necrophoresis, 1397
squares,
abscission, 68
Anthonomus grandis, 68
bolls, 68
stability of resistance, 1421, 1676
Chrysoperla carnea, 1676
Cry1Ac, 1421
fitness costs, 1421
Helicoverpa zea, 1421
heritability of resistance, 1421
organophosphates, 1676
pyrethroids, 1676
resistance, 1676
stand characteristics,
Dendroctonus, 1253
Ips, 1253
multiple regression models, 1253
Pinus ponderosa, 1253

Steinernema carpocapsae,
Heterorhabditis bacteriophora, 1533
multiple species approach, 1533
Otiorhynchus ligustici, 1533
Steinernema feltiae, 1533
Steinernema feltiae,
Heterorhabditis bacteriophora, 1533
multiple species approach, 1533
Otiorhynchus ligustici, 1533
Steinernema carpocapsae, 1533
Stelidota geminata,
insect movement, 850
integrated pest management, 850
stemborer, 50, 90
Busseola fusca, 90
Cotesia flavipes, 50
maize, 50, 90
nitrogen and potassium, 90
trap plants, 50
wild hosts, 50
yield loss, 90
sterile insect technique (SIT), 314, 1198
cactus moth, 314
circadian rhythm, 314
Cylas formicarius elegantulus, 1198
dispersal, 314
flight, 314
gamma radiation, 1198
mating behavior, 1198
sterile insect technique (SIT), 1198
sweetpotato weevil, 1198
sticky trap,
bait, 151
Blattella germanica, 151
trap bias, 151
trapping, 151
stink bugs, 1439, 1455
common smut, 1455
corn earworm, 1455
husk extension, 1455
insecticide, 1439
LC_{50}, 1439
maize weevil, 1455
sex-related response, 1439
storage, 42, 288
Cydia pomonella, 288
granulovirus, 288
microbial contaminants, 42
nucleopolyhedrovirus, 42
shelf life, 288
Spodoptera exigua, 42
temperature, 288
stored products, 605, 616
artificial diets, 616
booklice, 616
developmental rates, 605
life history, 605
marking, 605
psocids, 605
Psocoptera, 616
sex ratio, 616
stored-grain insects, 1011, 1022
barley, 1011
disinfestation, 1011
microwaves, 1011
mortality, 1011
spinosad, 1022
susceptibility differences, 1022
stored-product insects,
community ecology, 998
feed mills, 998
insect survey, 998
trapping, 998
straw mulch,
cultural management, 1317
habitat management, 1317
onions, 1317
Thrips tabaci, 1317
strawberry,
biological control, 251
entomopathogenic nematodes, 251
microbial control, 251
Striacosta albicosta,
corn pest, 404
monitoring, 404
pest management, 404
structure–activity relationship,
botanical insecticide, 1960
cinnamaldehyde, 1960
natural fumigant, 1960
Sitophilus oryzae, 1960
structure–activity relationship, 1960
sublethal dosages, 647, 1743
behavioral effects, 647
foraging behavior, 1743
honey bee, 1743
imidacloprid, 1743
orchard pollination, 647
side effects, 647
sublethal effects, 564, 784, 1805
antibiosis, 564
Apolygus lucorum, 1805
Brachiaria, 564
endosulfan, 1805
Megachile rotundata, 784
methodology, 784
Osmia lignaria, 784
pesticides, 784
resistance, 564
spittlebugs, 564
toxicity, 1805
Sucrocide,
Apis mellifera, 1057
desert conditions, 1057
oxalic acid, 1057
Varroa destructor, 1057
sugarcane, 17, 237, 334, 686
Chilo partellus, 686
climatic conditions, 334
COI, 686
Cotesia flavipes, 686
Delphacidae, 17
Dermolepida albohirtum, 334
Ethiopia, 686
Mexican rice borer, 237
movement, 237
pan evaporation, 334
Reoviridae, 17
rice, 237
Saccharum, 17
time series, 334
sugarcane borer,
Diatraea saccharalis, 1903
resistance, 1903
selection, 1903
tebufenozide, 1903
supplemental food,
Coccinellidae, 1792
Hippodamia variegata, 1792
marginal vegetation, 1792
Surround WP,
behavior, 118
oviposition, 118
particle film technology, 118
Rhagoletis mendax, 118
survival, 637, 777, 866, 989, 1798
Ceratitis capitata, 866
citrus peel, 866
development, 866, 1798
development thresholds, 1798
fecundity, 777
Galleria mellonella, 777
heat treatments, 989
host plant resistance, 866
humidity, 637
lysozyme, 777
Menacanthus stramineus, 637
modeling, 989
Ornithonyssus sylviarum, 637
reproduction, 1798
sodium tetraborate, 777
temperature, 637
thermal constants, 1798
Tribolium confusum, 989
survival time,
Agriotes obscurus, 375
fipronil, 375
insecticide toxicity, 375
wireworm, 375
susceptibility, 515, 1415
Chilo suppressalis, 1415
fipronil, 1415
imidacloprid, 515

metabolism, 1415
Nilaparvata lugens, 515
resistance, 515
risk assessment, 515
Sesamia inferens, 1415
susceptibility differences,
spinosad, 1022
stored-grain insects, 1022
sweep net sampling,
Oebalus pugnax, 216
Oryza sativa, 216
rice stink bug, 216
visual sampling, 216
sweetpotato,
Chaetocnema confinis, 1238
harvest date, 1238
planting date, 1238
soil temperature, 1238
sweetpotato weevil,
Cylas formicarius elegantulus, 1198
gamma radiation, 1198
mating behavior, 1198
sterile insect technique (SIT), 1198
symbiotic control,
Alcaligenes xylosoxidans, 1519
arthropod vector, 1519
paratransgenesis, 1519
Pierce's disease, 1519
Syrphidae,
aphid predators, 1526
conservation biocontrol, 1526
insectary crops, 1526
organic agriculture, 1526

T

table grape,
black widow spider, 1515
controlled atmosphere, 1515
postharvest pest control, 1515
ultralow oxygen, 1515
tasar silk worm,
eco-races, 1176
genetic variability, 1176
morphological traits, 1176
polymorphism, 1176
tebufenozide,
Diatraea saccharalis, 1903
resistance, 1903
selection, 1903
sugarcane borer, 1903
temperature, 288, 541, 637, 1557
Anarsia lineatella, 1557
bionomics, 1557
Cydia pomonella, 288
development, 541
developmental rate, 1557
granulovirus, 288
humidity, 637
Menacanthus stramineus, 637
model evaluation, 1557
Ornithonyssus sylviarum, 637
reproduction, 541
Russian wheat aphid, 541
shelf life, 288
storage, 288
survival, 637
Tenebrionidae,
Alphitobius diaperinus, 61
insect pest, 61
movement, 61
poultry litter, 61
Tephritidae,
invasive species, 750
Olea europaea, 750
olive fruit fly, 750
performance, 750
termite, 885, 1367
Coptotermes formosanus, 1367
dwellings, 1367
entomopathogenic fungus, 885
epizootic, 885
flooding, 1367
Metarhizium, 885
termite mortality,
neonicotinoids, 1881
soil penetration, 1881
treatment depth, 1881
termiticide, 902, 1360
horizontal transfer, 902
repellency, 1360
Reticulitermes flavipes, 902
Reticulitermes hesperus, 1360
soil treatments, 1360
Tetranychus urticae,
acaricide resistance, 1704
cross-resistance, 1704
hitchhiking effect, 1704
multiple resistance, 1704
Tettigoniidae,
Cicadellidae, 1779
clean stock program, 1779
Diaspididae, 1779
quarantine, 1779
thermal constants,
development, 1798
development thresholds, 1798
reproduction, 1798
survival, 1798
Thrips,
cropping-structure, 1685
insecticide-resistance, 1685
Thrips tabaci, 438, 1317, 1331
Allium cepa, 1331
cabbage, 438
cultural management, 1317
habitat management, 1317
insecticide, 438
onion thrips, 438, 1331
onions, 1317
plant resistance, 438
straw mulch, 1317
within-plant distribution, 1331
Thysanoptera,
acrinathrin, 499
fitness, 499
insecticide resistance, 499
spinosad, 499
time series,
climatic conditions, 334
Dermolepida albohirtum, 334
pan evaporation, 334
sugarcane, 334
timing of death,
Bactrocera dorsalis, 736
eclosion time, 736
heat exposure, 736
puparial morphology, 736
tobacco budworm, 1244, 1950
Bacillus thuringiensis, 1950
bollworm, 1244, 1950
cotton, 1950
integrated pest management, 1950
transgenic cotton, 1244
Vip3A, 1244
VipCot, 1244
tolerance,
Bt cotton, 959
economic threshold, 959
F_1 screen, 959
host plant resistance, 959
potato leafhopper, 959
resistance detection, 959
sap-feeding insect, 959
Tomarus subtropicus,
management, 1847
seasonal phenology, 1847
turfgrass, 1847
white grubs, 1847
tomato,
eggplant, 859
Leptinotarsa decemlineata, 859
potato, 859
solanaceous, 859
toxicity, 360, 1373, 1805
Apolygus lucorum, 1805
bifenthrin, 1373
endosulfan, 1805
fipronil, 1373
half-life, 1373
imidacloprid, 1373
insecticide, 360
silkworm, 360
sublethal effects, 1805
tracheal mites,
Acarapis woodi, 1756
Apis mellifera, 1756
genetic resistance, 1756
honey bees, 1756
transgenic cotton, 168, 1244
bollworm, 1244
Heliothis virescens, 168
insecticide, 168
resistance management, 168

tobacco budworm, 1244
Vip3A, 1244
VipCot, 1244
transmission,
Hemiptera, 1122
Pierce's disease, 1122
quantification, 1122
real-time polymerase chain reaction, 1122
transmission efficiency,
almond leaf scorch, 1105
grass sharpshooter, 1105
Pierce's disease (PD) of grapevines, 1105
population dynamics, 1105
trap, 126, 1838
attractant, 126
Ceratitis capitata, 126
Conotrachelus nenuphar, 1838
fruit fly, 126
lure, 126
material effect, 1838
monitoring, 1838
weather effect, 1838
trap bias,
bait, 151
Blattella germanica, 151
sticky trap, 151
trapping, 151
trap height,
emerald ash borer, 1831
prism trap, 1831
purple, 1831
visual cues, 1831
trap log,
Atimia confusa confusa, 430
Callidiellum rufipenne, 430
Callidium frigidum, 430
Semanotus ligneus ligneus, 430
trap plants,
Cotesia flavipes, 50
maize, 50
stemborer, 50
wild hosts, 50
trapping, 151, 399, 631, 998
bait, 151
Blattella germanica, 151
boll weevil, 399
community ecology, 998
dichlorvos, 399
ethylene glycol, 631
feed mills, 998
grandlure, 399
insect survey, 998
monitoring, 399
parasitoid, 631
Plodia interpunctella, 631
Sitophilus spp., 631
sticky trap, 151
stored-product insects, 998
trap bias, 151
treatment depth,
neonicotinoids, 1881
soil penetration, 1881
termite mortality, 1881
Tribolium confusum,
heat treatments, 989
modeling, 989
survival, 989
Trichogramma exiguum,
augmentation biological control, 421
microclimate, 421
predation, 421
Rhyacionia frustrana, 421
trichomes,
defoliation, 944
elm leaf beetle, 944
leaf toughness, 944
Ulmus spp., 944
Trichoplusia ni,
indoxacarb, 461
Pieris rapae, 461
Plutella xylostella, 461
spinosad, 461
trimedlure,
Ceratitis capitata, 1325
dispenser, 1325
fruit fly, 1325
pheromone, 1325
Triticum aestivum,
biotype, 569
Diuraphis noxia, 569
plant resistance, 569
turfgrass, 533, 1847, 1856
Blissus occiduus, 533
chinch bug, 533
granule formulation, 1856
management, 1847
mating disruption, 1856
oriental beetle, 1856
plant resistance, 533
seasonal phenology, 1847
soywax, 1856
Tomarus subtropicus, 1847
white grubs, 1847

U

Ulmus spp.,
defoliation, 944
elm leaf beetle, 944
leaf toughness, 944
trichomes, 944
ultralow oxygen,
black widow spider, 1515
controlled atmosphere, 1515
postharvest pest control, 1515
table grape, 1515
unicolonial,
ant baits, 1068
ant management, 1068
invasive species, 1068
little fire ant, 1068
UV-absorbing plastic,
Circulifer spp., 1337
Exitianus capicola, 1337
Orosius orientalis, 1337
physical control, 1337

V

Varroa destructor, 654, 668, 1057, 1075
Acarapis woodi, 668
Acarapis woodi, 654
Apis mellifera, 654, 668, 1057, 1075
desert conditions, 1057
Nosema spp., 668
oxalic acid, 1057
package honey bees, 668
PVC strips, 1075
residues, 1075
rotenone, 1075
Russian bees, 654
Sucrocide, 1057
varroa sensitive hygiene, 654
varroa mites,
acetic acid, 256
deformed wing virus, 256
formic acid, 256
fumigation, 256
varroa sensitive hygiene,
Acarapis woodi, 654
Apis mellifera, 654
Russian bees, 654
Varroa destructor, 654
vineyard, 699, 1287
Argentine ant, 699
deficit irrigation, 1287
grape, 1287
leafhoppers, 1287
Linepithema humile, 699
liquid baits, 699
Pseudococcidae, 699
water, 1287
Vip3A,
bollworm, 1244
tobacco budworm, 1244
transgenic cotton, 1244
VipCot, 1244
VipCot,
bollworm, 1244
tobacco budworm, 1244
transgenic cotton, 1244
Vip3A, 1244
visual cues,
emerald ash borer, 1831
prism trap, 1831
purple, 1831
trap height, 1831
visual sampling,
Oebalus pugnax, 216
Oryza sativa, 216
rice stink bug, 216
sweep net sampling, 216

W

water,
deficit irrigation, 1287
grape, 1287
leafhoppers, 1287
vineyard, 1287
weather effect,
Conotrachelus nenuphar, 1838
material effect, 1838
monitoring, 1838
traps, 1838
weed biological control,
Argopistes tsekooni, 1146
flea beetle, 1146
host specificity, 1146
Ligustrum sinense, 1146
weevil,
Conotrachelus nenuphar, 1302
integrated pest management, 1302
pheromone, 1302
plum curculio, 1302
western bean cutworm,
Bacillus thuringiensis corn, 555
corn hybrids, 555
insect injury, 555
western corn rootworm,
Cuphea, 797
Diabrotica virgifera, 797
white grubs, 1158, 1624, 1847
Cynodon, 1624
cytochrome c oxidase I, 1624
management, 1847
nursery, 1158
Phyllophaga, 1624
Popillia japonica, 1158
quarantine, 1158
scarab, 1158
scarab flight periods, 1624
seasonal phenology, 1847
Tomarus subtropicus, 1847
turfgrass, 1847
whitefly immature stages,
Bemisia argentifolii, 174
imidacloprid-resistant whiteflies, 174
Oberon, 174
Q Biotype, 174
Widestrike,
Bollgard II, 1594
Bollworm, 1594
Helicoverpa, 1594
integrated pest management, 1594
wild hosts,
Cotesia flavipes, 50
maize, 50
stemborer, 50
trap plants, 50
winter colony growth,
brood pheromone, 1749
honey bee, 1749
wireworm, 365, 375, 1568
Agriotes obscurus, 375
Elateridae, 365
fipronil, 375
insecticide toxicity, 365, 375
mating disruption, 1568
mating rate, 1568
morbidity, 365
neonicotinoids, 365
sex pheromone, 1568
survival time, 375
within-canopy distribution,
Cydia pomonella, 81
fruit phenology, 81
larval infestation, 81
monitoring, 81
within-plant distribution,
Allium cepa, 1331
onion thrips, 1331
Thrips tabaci, 1331

X

Xenopus oocyte,
abamectin, 1888
functional expression, 1888
GABA receptor, 1888
Plutella xylostella, 1888
Xylella fastidiosa,
epidemiology, 1041
Homalodisca vitripennis, 1041
Pierce's disease, 1041
xylose,
bait, 873
carbohydrates, 873
feeding, 873
[^{14}C]hexaflumuron, 873

Y

yam chips,
Benin, 1967
cassava chips, 1967
infestation, 1967
Prostephanus truncatus, 1967
yellow sticky trap,
glassy-winged sharpshooter, 226
orange trees, 226
pest management, 226
sex ratio, 226
YHD2,
Cry1Ac, 1406
Heliothis virescens, 1406
insecticide resistance management, 1406
single-pair family, 1406
yield loss,
Busseola fusca, 90
maize, 90
nitrogen and potassium, 90
stemborer, 90
Yucatán,
bees, 132
crop pollination, 132
greenhouse, 132

Z

Zea mays,
Diabrotica virgifera virgifera, 341
fatty acid, 341
feeding stimulants, 341
insect behavior, 341